Combination Theorems in Groups, Geometry and Dynamics

Mahan Mj and Sabyasachi Mukherjee

Abstract The aim of this article is to give a survey of combination theorems occurring in hyperbolic geometry, geometric group theory and complex dynamics, with a particular focus on Thurston's contribution and influence in the field.

Key words: Kleinian group, hyperbolic group, flaring, holomorphic mating, simultaneous uniformization, double limit theorem

Sabyasachi Mukherjee

Mahan Mj

School of Mathematics Tata Institute of Fundamental Research, Mumbai-400005 India, e-mail: mahan@math.tifr.res.in

School of Mathematics Tata Institute of Fundamental Research, Mumbai-400005 India, e-mail: sabya@math.tifr.res.in

Combi Mahan	ination T	heorems in Groups, Geometry and Dynamics	1
1	1 Introduction		
2	Klei	Klein-Maskit combination for Kleinian groups	
3	3 Simultaneous uniformization and Quasi-Fuchsian Groups		7
-	3.1	Topologies on space of representations	7
	3.2	Simultaneous uniformization	8
	3.3	Geodesic Laminations	10
4	Thu	Thurston's combination theorem for Haken manifolds	
	4.1	Non-fibered Haken 3-manifolds	12
	4.2	The double limit theorem	13
5	Com	Combination theorems in geometric group theory: hyperbolic	
	grou	ps	14
	5.1	Trees of spaces	15
	5.2	Metric Bundles	17
	5.3	Relatively hyperbolic combination theorems	21
	5.4	Effective quasiconvexity and flaring	23
6	Com	ombination theorems in geometric group theory: cubulations 25	
7	Holo	Holomorphic dynamics and polynomial mating	
	7.1	Historical comments	27
	7.2	Mating of polynomials	28
8	Com	Combining rational maps and Kleinian groups	
	8.1	Mating anti-polynomials with reflection groups	32
	8.2	Mating polynomials with Kleinian groups	41
Re	References		
Re	eferences		47

1 Introduction

The aim of this survey is to give an eclectic account of combination theorems in hyperbolic geometry, geometric group theory and complex dynamics. Thurston's contribution and influence in the theme is pervasive, and we will only be able to touch upon some of these aspects. The hope in writing this survey is therefore only to whet the appetite of the reader and provide some references to more detailed articles and surveys.

Combination theorems have a long history, going back to Klein's paper from 1883 [Kle83]. A major subsequent development in terms of combination theorems for Kleinian groups is due to Maskit [Mas65b, Mas68, Mas71, Mas93, Mas80]. In the 1970's and 80's, a phase transition occurred in the theory with the advent of Thurston and his proof of hyperbolization of atoroidal Haken manifolds [Thu86a, Thu86b, Thu86c, Ota01, Ota98, Kap01b]. Thurston's work has had a deep and profound influence on hyperbolic geometry ever since, and has provided a template for related developments in geometric group theory and complex dynamics. In geometric group theory, Bestvina and Feighn [BF92] isolated the coarse geometric features of Thurston's combination theorem and proved a highly influential combination theorem for Gromov-hyperbolic groups [Gro85], spawning considerable activity and several generalizations [Dah03, Ali05, MR08, MS12, Gau16]. Finally, in a relatively recent major development leading to a resolution of Thurston's virtual Haken conjecture by Agol and Wise [Ago13, Wis21], Haglund and Wise [HW12] proved a combination theorem for virtually special cubulable hyperbolic groups [HW08].

In a closely related theme, Fatou had already observed similarities between limit sets of Kleinian groups and Julia sets of rational maps in the 1920s [Fat29, p. 22]: 'L'analogie remarqueé entre les ensembles de points limites des groupes Kleineens et ceux qui sont constitués par les frontières des régions de convergence des itérées d'une fonction rationnelle ne parait d'ailleurs pas fortuite et il serait probablement possible d'en faire la syntése dans une théorie générale des groupes discontinus des substitutions algrébriques.' The theory of mating, developed by Douady and Hubbard [Dou83] extends the notion of a combination theorem from the world of Kleinian groups to that of complex dynamics. This theme too bears the tell-tale stamp of Thurston. In fact, Thurston's topological characterization of rational maps is an invaluable tool in constructing such matings [DH93, Shi00].

2 Klein-Maskit combination for Kleinian groups

A discrete subgroup Γ of $PSL_2(\mathbb{C})$ is called a *Kleinian group*. The *limit set* of the Kleinian group Γ , denoted by Λ_{Γ} , is the collection of accumulation points of a Γ -orbit $\Gamma \cdot z$ for some $z \in \hat{\mathbb{C}}$. Λ_{Γ} is independent of z. It may be thought of as the locus of chaotic dynamics of Γ on $\hat{\mathbb{C}}$, i.e. for Γ non-elementary and any $z \in \Lambda_{\Gamma}$, $\Gamma \cdot z$ is dense in Λ_{Γ} . We shall identify the Riemann sphere $\hat{\mathbb{C}}$ with the sphere at infinity \mathbb{S}^2 of \mathbb{H}^3 . The complement of the limit set $\hat{\mathbb{C}} \setminus \Lambda_{\Gamma}$ is called the domain of discontinuity $\Omega(\Gamma)$ of

 Γ . If the Kleinian group Γ is torsion-free, it acts freely and properly discontinuously on $\Omega(\Gamma)$ with a Riemann surface quotient.

Definition 1 A set *D* is called a *partial fundamental domain* for Γ , if

1. $D \neq \emptyset$,

2. $D \subset \Omega(\Gamma)$, and

3. $g(D) \cap D = \emptyset$, for all $g \in \Gamma$, $g \neq 1$.

If, further, $\bigcup_{g \in \Gamma} g.D = \Omega(\Gamma)$, then *D* is called a *fundamental domain* for Γ .

The story of combination theorems starts with the following theorem of Klein:

Theorem 1 (Klein combination theorem) [*Kle83*] Let Γ_1, Γ_2 be Kleinian groups with fundamental domains D_1, D_2 respectively. Assume that the interior of D_1 (resp. D_2) contains the boundary and exterior of D_2 (resp. D_1). Then the group Γ generated by Γ_1, Γ_2 is Kleinian, and $D = D_1 \cap D_2$ is a fundamental domain for Γ .

In the 1960's, Maskit started working on extending the Klein combination Theorem 1 to a more general setup. Maskit's work on combination theorems for Kleinian groups started with the following.

Theorem 2 (Klein-Maskit combination theorem for free product with amalgamation) [*Mas*65*a*] Let Γ_1, Γ_2 be Kleinian groups with domains of discontinuity Ω_1, Ω_2 respectively. Let $H = \Gamma_1 \cap \Gamma_2$. Let D_1, D_2, Δ be partial fundamental domains for Γ_1, Γ_2, H respectively. For i = 1, 2, set $E_i = H.D_i$. Denote the interior of $D = E_1 \cap E_2 \cap \Delta$ by D'. If

 $1. D' \neq \emptyset,$ $2. E_1 \cup E_2 = \Omega_1 \cup \Omega_2.$

Then the group Γ generated by Γ_1, Γ_2 is Kleinian, D' is a partial fundamental domain for Γ , $\Gamma = \Gamma_1 *_H \Gamma_2$ is the free product amalgamation of Γ_1, Γ_2 along H. Further, $gD \cap D = \emptyset$, for all $g \in \Gamma$, $g \neq 1$.

In [Mas65b], Maskit strengthened the above theorem by determining precisely a fundamental domain for the group.

Theorem 3 (Klein-Maskit combination theorem for free product with amalgamation) [*Mas*65*b*] Let Γ_1, Γ_2 be Kleinian groups with domains of discontinuity Ω_1, Ω_2 respectively. Let $H \subset \Gamma_1 \cap \Gamma_2$. such that *H* is either cyclic or consists only of the identity. Let D_1, D_2, Δ be fundamental domains for Γ_1, Γ_2, H respectively. For i = 1, 2, set $E_i = H.D_i$. Denote the interior of $D = E_1 \cap E_2 \cap \Delta$ by *D'*. Suppose $E_1 \cup E_2 = \Omega(H)$ and that $D' \neq \emptyset$. Assume further that there is a simple closed curve γ , contained in $int(E_1 \cup E_2) \cup \Lambda_H$ such that γ is invariant under *H*; the closure of $\gamma \cap \Delta$ is contained in $int(E_1 \cap E_2)$ and γ separates both $E_1 \setminus E_2$ and $E_2 \setminus E_1$. Then the group Γ generated by Γ_1, Γ_2 is Kleinian, $\Gamma = \Gamma_1 *_H \Gamma_2$, and *D* is a fundamental domain for *G*. Subsequently, in [Mas68, Mas71], Maskit upgraded Theorem 3 to the following. We start with two Kleinian groups Γ_1, Γ_2 with $H \subset \Gamma_1 \cap \Gamma_2$, where $H \neq \Gamma_1, \Gamma_2$. We are also given a simple closed curve γ dividing the Riemann sphere $\hat{\mathbb{C}}$ into two closed topological discs, B_1 and B_2 , where B_i is precisely invariant under H in Γ_i . More precisely, B_i is H-invariant, and if $g \in \Gamma_i \setminus H$, then $g(B_i) \cap B_i = \emptyset$. Then $\Gamma = \langle \Gamma_1, \Gamma_2 \rangle$, the group generated by Γ_1 and Γ_2 is also a Kleinian group. What really needs to be proved in all these cases is the discreteness of Γ .

In all these cases, Maskit shows that $\Gamma = \Gamma_1 *_H \Gamma_2$, i.e. $\Gamma = \langle \Gamma_1, \Gamma_2 \rangle$ equals the free product with amalgamation of Γ_1, Γ_2 along *H*. Further, by carefully choosing fundamental domains for Γ_1, Γ_2 one can ensure that their intersection will be a fundamental domain for Γ . Thus, the basic hypothesis guaranteeing discreteness of Γ can be summarized as follows:

- 1. The disks B_1 and B_2 are both invariant under H.
- 2. The $(\Gamma_1 \setminus H)$ -translates of B_1 are disjoint disks in B_2 .
- 3. The $(\Gamma_2 \setminus H)$ -translates of B_2 are disjoint disks in B_1 .

There is also a version of the Klein-Maskit combination theorem for HNN extensions. We are given a single group Γ_0 , with two subgroups H_1 and H_2 , two closed disks B_1 and B_2 , which have disjoint projections to $\Omega(\Gamma_0)/\Gamma_0$, where

- 1. H_i preserves B_i ,
- 2. there exists a Möbius transformation h mapping the outside of B_1 onto the inside of B_2 and conjugating H_1 to H_2 .

Maskit then shows that $\Gamma = \langle \Gamma_0, h \rangle$ is a Kleinian group. Also $\Gamma = \Gamma_0 *_H$ is the HNN-extension of G_0 along H, where the two inclusions of H map to H_1, H_2 and h is the stable letter conjugating one to the other. Further, by carefully choosing a fundamental domain D for Γ_0 , one can ensure that $D \setminus (B_1 \cup B_2)$ is a fundamental domain for Γ .

Maskit weakens the hypotheses further in [Mas93], allowing translates of the closed disks B_1, B_2 to have common boundary points. However, in [Mas93] he requires that such points of intersection also be ordinary points of our original group. In [Mas93], it is also shown that Γ is geometrically finite if and only if the original groups are so. The basic topological tool used in the proof is a Jordan curve γ in $\hat{\mathbb{C}}$ and its translates under a Kleinian group. The standard hypothesis in these papers is the 'almost disjointness' of γ from all its translates. More precisely, if $g(\gamma) \cap \gamma \neq \emptyset$, then it is required that $g(\gamma)$ is entirely contained in the closure of one of the open disks bounded by γ . Thus, a substantial amount of the technical difficulty in [Mas71, Mas93] comes from controlling the points of intersection $g(\gamma) \cap \gamma$.

To conclude this section, we refer the reader to

- Work of Li-Ohshika-Wang [LOW09, LOW15] for generalizations of the Klein-Maskit combination theorems to higher dimensions.
- 2. Work of Dey, Kapovich and Leeb [DKL14] for a combination theorem for Anosov subgroups, a natural class of discrete subgroups of higher rank Lie groups that generalizes convex cocompact subgroups of $PSL_2(\mathbb{C})$.

3 Simultaneous uniformization and Quasi-Fuchsian Groups

The aim of this section is to give a brief account of Bers' simultaneous uniformization theorem. The reason is 2-fold. First, it provides the context for Thurston's double limit theorem in Section 4.2. Second, it is the Kleinian group analog for the Douady-Hubbard mating construction [Dou83], and more generally the original motivation for the mating constructions in Section 8.

Fix a surface *S*. The collection of all representations $\rho : \pi_1(S) \to PSL_2(\mathbb{C})$ up to conjugacy (in $PSL_2(\mathbb{C})$) is called the *character variety* and is represented as $\mathcal{R}(S)$. We note in passing that the appropriate quotient by $PSL_2(\mathbb{C})$ of the space of all representations $\rho : \pi_1(S) \to PSL_2(\mathbb{C})$ is the GIT quotient. This is needed in order to obtain the structure of a variety on $\mathcal{R}(S)$.

3.1 Topologies on space of representations

For future reference, we summarize here a natural collection of topologies on the space of *discrete faithful* ρ : $\pi_1(S) \rightarrow PSL_2(\mathbb{C})$. The *algebraic topology* is the topology of pointwise convergence on elements of $\pi_1(S)$:

Definition 2 We shall say that a sequence of representations $\rho_n : \pi_1(S) \to PSL_2(\mathbb{C})$ converges *algebraically* to $\rho_{\infty} : \pi_1(S) \to PSL_2(\mathbb{C})$ if for all $g \in \pi_1(S)$, $\rho_n(g) \to \rho_{\infty}(g)$ in $PSL_2(\mathbb{C})$.

The collection of conjugacy classes of discrete faithful representations of $\pi_1(S)$ into $PSL_2(\mathbb{C})$ equipped with the algebraic topology is denoted as AH(S). Thus, $AH(S) \subset \mathcal{R}(S)$ comes naturally equipped with a complex analytic structure. The space of discrete faithful representations of $\pi_1(S)$ into $PSL_2(\mathbb{R})$ equipped with the algebraic topology is precisely the Teichmüller space. Thus, the Teichmüller space sits 'diagonally' in AH(S).

For analyzing convergence from a geometric point of view, the natural topology is the *geometric topology*, or equivalently, the Gromov-Hausdorff topology.

Definition 3 Let $\rho_n : \Gamma \to PSL_2(\mathbb{C})$ be a sequence of discrete, faithful representations of a finitely generated, torsion-free, nonabelian group Γ . Thus, $\rho_n(\Gamma)$ is a sequence of closed subsets of $PSL_2(\mathbb{C})$. If $G \subset PSL_2(\mathbb{C})$ is a closed subgroup, such that $\rho_n(\Gamma)$ converges to G in the Gromov-Hausdorff topology, then $\rho_n(\Gamma)$ is said to *converge geometrically* to G, and G is called the *geometric limit*.

Definition 4 G_n converges *strongly* to G if G_n converges to G both algebraically and geometrically.

3.2 Simultaneous uniformization

Definition 5 Let $\rho : \pi_1(S) \to PSL_2(\mathbb{C})$ be a discrete faithful representation such that the limit set of $G = \rho(\pi_1(S))$ is a topological circle in \mathbb{S}^2 . Then *G* is said to be *quasiFuchsian*. The collection of conjugacy classes of quasi-Fuchsian representations is denoted as QF(S).

Note that QF(S) is contained in AH(S) and hence inherits a complex analytic structure. The domain of discontinuity Ω of a quasi-Fuchsian *G* consists of two open invariant disks Ω_1, Ω_2 . Hence the quotient Ω/G is the disjoint union $\Omega_1/G \sqcup \Omega_2/G$. Hence we have a map $\tau : QF(S) \to Teich(S) \times Teich(S)$, where Teich(S) denotes the Teichmüller space of *S*. The **Bers simultaneous Uniformization Theorem** asserts:

Theorem 4 [Ber60, Ber61] τ : $QF(S) \rightarrow Teich(S) \times Teich(S)$ is a homeomorphism.

Hence, given any two conformal structures T_1, T_2 on a surface, there is a unique discrete quasi-Fuchsian *G* whose limit set Λ_G is topologically a circle, and the quotient of whose domain of discontinuity is $T_1 \sqcup T_2$. See Figure 1 below [Kab16], where the inside and the outside of the Jordan curve correspond to Ω_1, Ω_2 .



Fig. 1 Quasi-Fuchsian group limit set

We refer to [Hub06] for a proof of Theorem 4 and summarize the main ideas here. Theorem 4 is essentially complex analytic in nature and goes back to an understanding of the Beltrami partial differential equation due to Morrey. Let K_S denote the canonical bundle of the Riemann surface S (if S has punctures as a hyperbolic surface, we regard them as marked points in the complex analytic category).

Definition 6 A *Beltrami differential* on *S* is an L^{∞} section of $K_S^{-1} \otimes \overline{K_S}$, where $\overline{K_S}$ denotes the complex conjugate of K_S . The space of Beltrami differentials on *S* will be denoted as $\mathcal{D}_b(S)$

The local expression for an element of $\mathcal{D}_b(S)$ in a complex analytic chart $U \subset S$ is thus given by $\mu \frac{d\overline{z}}{dz}$, where $\mu \in L^{\infty}(U)$ is called a *Beltrami coefficient*.

Definition 7 A *quasiconformal map* between two Riemann surfaces S_1 and S_2 is a homeomorphism $\phi : S_1 \to S_2$ having locally square-integrable weak partial derivatives such that

$$\mu = \frac{\phi_{\overline{z}}}{\phi_z}$$

satisfies $||\mu||_{\infty} < 1$. Here, μ is called the *Beltrami coefficient* of ϕ .

The first major ingredient in the proof of Theorem 4 is the Measurable Riemann mapping theorem. As pointed out by Hubbard in [Hub06, p. 149], the Beltrami coefficient μ really represents an almost-complex structure on U and the Measurable Riemann mapping theorem (due to Ahlfors-Bers-Morrey) below ensures its integrability to a complex structure.

Theorem 5 (Measurable Riemann Mapping Theorem) [Hub06, Theorem 4.6.1] **Existence of quasiconformal maps:** Let $U \subset \mathbb{C}$ be open. Let $\mu \in L^{\infty}(U)$ satisfying $||\mu||_{\infty} < 1$. Then there exists a quasiconformal mapping $f : U \to \mathbb{C}$ solving the Beltrami equation

$$\frac{\partial f}{\partial \overline{z}} = \mu \frac{\partial f}{\partial z}.$$

Uniqueness of quasiconformal maps: *If g is another quasiconformal solution to the Beltrami equation above, then there exists a univalent analytic function* ϕ : $f(U) \rightarrow \mathbb{C}$ such that $g = \phi \circ f$.

The rest of this brief account of Theorem 4 follows [GS20] which captures the relevant conformal geometry. Recall that we have fixed a base Riemann surface *S*. Let $\Gamma < PSL_2(\mathbb{R})$ be a (base) Fuchsian group uniformizing *S*. Let *S'* be an arbitrary point in the Teichmüller space of the underlying topological surface. Theorem 4 then associates to *S'* a quasiconformal map $\Phi : \hat{\mathbb{C}} \to \hat{\mathbb{C}}$ fixing the three points 0, 1, ∞, and conjugating the action of Γ to that of a Kleinian group $\Gamma(S, S')$, such that

- 1. Φ is conformal on the lower half-plane.
- 2. $\Gamma(S, S')$ leaves invariant the images of the lower and upper half-planes.
- 3. The quotient of the lower half-plane by $\Gamma(S, S')$ is *S*.
- 4. The quotient of the upper half-plane by $\Gamma(S, S')$ is S'.

To prove the existence of a Φ as above, we first note that the Teichmüller space can be identified (via Theorem 5) with Beltrami differentials on *S* with norm bounded by one. Let μ be the Beltrami differential on *S* corresponding to *S'*. Next. lift μ to the upper half plane. Extend to a Beltrami coefficient μ_0 on $\hat{\mathbb{C}}$ by defining it to be identically zero on the lower half plane. The map Φ above is then given by the normalized solution to the Beltrami equation with the Beltrami coefficient μ_0 . Invariance of μ_0 under Γ ensures the existence of an isomorphism $\rho : \Gamma \to \Gamma(S, S')$ to the desired Kleinian group $\Gamma(S, S')$ such that

$$\Phi \circ g = \rho(g) \circ \Phi,$$

for all $g \in \Gamma$.

3.3 Geodesic Laminations

We turn now to the hyperbolic geometry of quasi-Fuchsian groups.

Definition 8 A geodesic lamination on a hyperbolic surface is a foliation of a closed subset with geodesics.

A geodesic lamination on a surface may further be equipped with a transverse (positive) measure to obtain a *measured lamination*. The space of measured (geodesic) laminations on S is then a positive cone in a vector space and is denoted as $\mathcal{ML}(S)$. It can be projectivized to obtain the space of projectivized measured laminations $\mathcal{PML}(S)$. It was shown by Thurston [FLP79] that

Theorem 6 $\mathcal{PML}(S)$ is homeomorphic to a sphere and can be adjoined to Teich(S) compactifying the latter to a closed ball.

Definition 9 [Thu80][Definition 8.8.1] A *pleated surface* in a hyperbolic threemanifold N is a complete hyperbolic surface S of finite area, together with an isometric map $f : S \to N$ such that every $x \in S$ is in the interior of some geodesic segment which is mapped by f to a straight line segment. Also, f must take every cusp of S to a cusp of N

The pleating locus of the pleated surface $f : S \to M$ is the set $\gamma \subset S$ consisting of those points in the pleated surface which are in the interior of unique line segments mapped to line segments.

Proposition [Thu80][Proposition 8.8.2] The pleating locus γ is a geodesic lamination on S. The map f is totally geodesic in the complement of γ .

The geometry of quasi-Fuchsian groups and their relationship with geodesic laminations arises out of the geometry of the convex core that we now describe.

Definition 10 Let Γ be an infinite Kleinian group and let $\Lambda \subset \hat{\mathbb{C}}$ denote its limit set. The *convex hull* of Λ is the smallest non-empty closed convex subset of \mathbb{H}^3 whose set of accumulation points in $\hat{\mathbb{C}}$ equals Λ . We denote the convex hull of Λ by $CH(\Lambda)$.

The convex hull $CH(\Lambda)$ of a Kleinian group Γ is invariant under Γ . The quotient $CH(\Lambda)/\Gamma \subset \mathbb{H}^3/\Gamma$ is called the *convex core* of the hyperbolic 3-manifold $M = \mathbb{H}^3/\Gamma$.

For a quasi-Fuchsian group $\Gamma = \rho(\pi_1(S))$, the convex core is homeomorphic to a product $S \times [a, b]$ (the Fuchsian case corresponds to a = b). The hyperbolic distance between $S \times \{a\}$ and $S \times \{b\}$ is a measure of the geometric complexity of Γ . In [Thu80][Ch. 8], Thurston further shows:

Proposition Let *M* be a complete hyperbolic 3-manifold corresponding to a quasi-Fuchsian group Γ , and let CC(M) denote its convex core. Then each component of the convex core boundary $\partial CC(M)$ is a pleated surface.

4 Thurston's combination theorem for Haken manifolds

The material in this section provides the core inspiration for most combination theorems that came subsequently.

Definition 11 [Hem04] A properly embedded surface $(F, \partial F) \subset (M, \partial M)$ in a 3manifold M with boundary ∂M (possibly empty) is said to be *incompressible* if the inclusion map $i : (F, \partial F) \subset (M, \partial M)$ induces an injective homomorphism of fundamental groups $i_* : \pi_1(F) \to \pi_1(M)$. Further, we require that for every boundary component γ of ∂F , $i_* : \pi_1(\gamma) \to \pi_1(\partial M)$ is injective. (The second condition is automatic when F is not a disk.)

An embedded incompressible surface $(F, \partial F) \subset (M, \partial M)$ is said to be *boundary parallel* if *F* can be isotoped into ∂M keeping $\partial F \subset \partial M$ fixed.

A compact 3-manifold M (possibly with boundary ∂M) is said to be *Haken* if

- 1. $\pi_2(M) = 0$.
- 2. There exists an embedded incompressible surface $(F, \partial F)$ that is not boundary parallel.

M is said to be atoroidal if $\pi_1(M)$ contains no $\mathbb{Z} \oplus \mathbb{Z}$ subgroups. *M* is said to be acylindrical if any embedded incompressible annulus in $(M, \partial M)$ is boundary parallel.

We summarize Thurston's celebrated hyperbolization theorem now and then give a brief account of the ingredients that go into the proof.

Theorem 9 [*Thu86a, Thu86b, Thu86c*] Let *M* be a compact atoroidal Haken 3manifold. Then *M* is hyperbolic.

There is a version of Theorem 9 for 3-manifolds with torus boundary components also. But, in the interests of exposition, we shall largely focus on the compact atoroidal case. The proof of Theorem 9 breaks into two principal pieces:

- 1. *M* is compact atoroidal Haken and does not fiber over \mathbb{S}^1 . This case will be described in Section 4.1.
- 2. M fibers over the circle with fiber F. This case will be described in Section 4.2.

4.1 Non-fibered Haken 3-manifolds

There are a number of detailed expositions for the compact atoroidal Haken non-fibered case and we point out [Ota98, Kap01b, McM90] in particular.

It is a fundamental fact of 3-manifold topology [Hem04, Chapter 13] that any Haken manifold admits a Haken hierarchy. Cutting $(M, \partial M)$ open along $(F, \partial F)$ gives us a new (possibly disconnected) atoroidal 3-manifold with boundary. The cut open manifold is automatically Haken, and we can proceed inductively. At the last stage, we are left with a finite collection of balls, and these are clearly hyperbolic.

Thus, in order to prove Theorem 9 in the non-fibered case, an essential step is the following:

Theorem 10 [Ota98] Let M_1 be an acylindrical atoroidal 3-manifold with nonempty incompressible boundary ∂M_1 such that its interior admits a complete hyperbolic metric. Let $\tau : \partial M_1 \to \partial M_1$ be an orientation-reversing involution. Then the interior of $M = M_1/\tau$ admits a complete hyperbolic metric.

To prove Theorem 10, a first tool is the following generalization of Theorem 4:

Theorem 11 Let M_1 be a compact 3-manifold with boundary such that

1. The interior of M_1 admits a complete hyperbolic metric.

2. No component of ∂M_1 is homeomorphic to a torus or a sphere.

Then the space of complete hyperbolic metrics on M_1 is given by $Teich(\partial M_1)$.

Let $\partial M_1 = \bigsqcup_i \Sigma_i$, where each Σ_i is a surface of genus greater than one. Then $Teich(\partial M_1) = \prod_i Teich(\Sigma_i)$. Fix a complete hyperbolic structure on M_1 (the existence of such a structure is guaranteed by the hypothesis of Theorem 11). This is equivalent to a discrete faithful representation $\rho : \pi_1(M_1) \to PSL_2(\mathbb{C})$. Let $\Gamma = \rho(\pi_1(M_1))$. Then each $\Sigma_i \subset \partial M_1$ gives (via inclusion) a conjugacy class of quasi-Fuchsian subgroups of Γ . Thus the involution τ of Theorem 10 induces a map

$$\sigma: Teich(\partial M_1) \to Teich(\partial M_1).$$

The map σ is called the *skinning map*. The existence of a complete hyperbolic structure on $M = M_1/\tau$ is equivalent to the existence of a fixed point of the skinning map σ , as such a fixed point ensures an isometric gluing. Thurston's fixed point theorem can now be stated as the following: If M is atoroidal then σ has a fixed point. It now follows from the Klein-Maskit combination theorem (Section 2) that M admits a complete hyperbolic structure. The acylindricity hypothesis of Theorem 10 guarantees that M is atoroidal, completing an outline of the proof of Theorem 10.

An effective method of proving the existence of a fixed point of the skinning map σ was taken by McMullen in [McM90]. As mentioned in [Ota98] Hubbard had observed that the analytical formula for the coderivative of the skinning map relates it to the Theta operator in Teichmüller theory. McMullen studies the fixed point problem via $||D\sigma||$, the norm of the derivative of the skinning map. He reproves

Theorem 10 by showing that if M_1 is acylindrical, then there exists c < 1 such that $||D\sigma|| < c$ guaranteeing a solution to the gluing problem.

Both Thurston's fixed-point theorem and McMullen's estimates in [McM90] are easiest to state when M_1 is acylindrical. However, both approaches can be refined to conclude hyperbolicity of M so long as M_1 is not of the form $S \times I$ and τ glues $S \times \{0\}$ to $S \times \{1\}$. The excluded case is that of 3-manifolds fibering over the circle and involves a completely different approach that we describe now.

4.2 The double limit theorem

We shall follow [Ota01] to give an outline of the steps involved in the hyperbolization of 3-manifolds fibering over the circle. Recall (Theorem 6) that the space of projectivized measured laminations $\mathcal{PML}(S)$ compactifies Teich(S). Thurston's double limit theorem may be thought of as an extension of the simultaneous uniformization Theorem 4 to the case where the pair (τ_1, τ_2) of Riemann surfaces in $Teich(S) \times Teich(S)$ is replaced by a pair $(\ell_1, \ell_2) \in \overline{Teich(S)} \times \overline{Teich(S)}$, where $\overline{Teich(S)} = Teich(S) \cup \mathcal{PML}(S)$ denotes the Thurston compactification of Teich(S)as in Theorem 6.

Dual to any measured lamination $\ell \in \mathcal{ML}(S)$ there is an action of $\pi_1(S)$ on an \mathbb{R} -tree. An \mathbb{R} -tree is a geodesic metric space such that any two distinct points are joined by a unique arc isometric to an interval in \mathbb{R} . We refer to [Bes02] for an expository account of group actions on \mathbb{R} -trees and convergence of Γ -spaces, and mention only the following theorem. Fix a group Γ . A triple (X, o, ρ) is called a based Γ -space if $o \in X$ is a base-point, and Γ acts on X via a homomorphism $\rho : \Gamma \rightarrow Isom(X)$ from Γ to the isometry group Isom(X) of X.

Theorem 12 [Bes02][Theorem 3.3] Let (X_i, o_i, ρ_i) be a convergent sequence of based Γ -spaces such that

1. Each X_i is δ hyperbolic, for some $\delta \geq 0$.

2. there exists $g \in \Gamma$ such that the sequence $d_i = d_{X_i}(o_i, \rho_i(g)(o_i))$ is unbounded.

Then there is a based \mathbb{R} -tree (T, o) and an isometric action $\rho : \Gamma \to Isom(T)$ such that $(X_i, o_i, \rho_i) \to (T, o, \rho)$.

Further, the (pseudo)metric on the \mathbb{R} -tree T is obtained as the limit of pseudometrics $\frac{d_{(X_i,o_i,\rho_i)}}{d_i}$.

Finally, we shall need the following theorem of Skora [Sko96] on the structure of groups admitting small actions on \mathbb{R} -trees.

Theorem 13 [Sko96] Let S be a finite area hyperbolic surface. Suppose $\pi_1(S)$ acts non-trivially on an \mathbb{R} -tree T, such that for every cusp \mathbb{P} of S, $\pi_1(\mathbb{P})$ fixes a point in T. Then the stabilizer of each non-degenerate arc of T contains no free subgroup of rank 2 if and only if the action is dual to an element of $\mathcal{ML}(S)$.

An action of $\pi_1(S)$ on an \mathbb{R} -tree T such that the stabilizer of each non-degenerate arc of T contains no free subgroup of rank 2 is called a *small action*. Morgan and Shalen [MS84, MS88a, MS88b] constructed a compactification of the variety $\mathcal{R}(S)$ by the space of projectivized length functions arising from small actions of $\pi_1(S)$ on \mathbb{R} -trees. Skora's theorem 13 allows us to replace $\mathcal{PML}(S)$ in Theorem 6 by such length functions.

With this background in place we return to an outline of Thurston's double limit theorem [Thu86b] following Otal [Ota01]. Let $(\tau_i^+, \tau_i^-) \in Teich(S) \times Teich(S)$ be a sequence of points converging to $(\ell_+, \ell_-) \in \overline{Teich(S)} \times \overline{Teich(S)}$. By Theorem 4, we can identify $Teich(S) \times Teich(S)$ with QF(S) and hence assume that $(\tau_i^+, \tau_i^-) \in QF(S)$. For convenience of exposition, we assume that ℓ_+, ℓ_- are both in $\mathcal{PML}(S)$ (a similar statement holds if only one of ℓ_+, ℓ_- lies in $\mathcal{PML}(S)$). Assume further that ℓ_+, ℓ_- fill S, i.e. each component of $S \setminus (\ell_+ \cup \ell_-)$ is either simply connected or else is topologically a punctured disk. Let $\rho_i : \pi(S) \to PSL_2(\mathbb{C})$ be the quasi-Fuchsian representation corresponding to $(\tau_i^+, \tau_i^-) \in QF(S)$ and let $\Gamma_i = \rho_i(\pi(S))$. Thurston's double limit theorem now says:

Theorem 14 [*Thu86b*] Under the above assumptions, there exists a Kleinian group Γ such that Γ_i converges to Γ in AH(S).

We sketch Otal's proof following [Ota01] and argue by contradiction. If Γ_i diverges, then Theorem 12 shows that there is a limiting small action of $\pi_1(S)$ on an \mathbb{R} -tree *T*. By Theorem 13 such a small action is dual to a measured lamination ℓ on *S*.

It is then shown in [Ota01] that any measured lamination that intersects ℓ essentially is realizable in *T*. Hence at least one of ℓ_+ and ℓ_- must be realizable in *T*, since the two together fill *S*. Without loss of generality, suppose ℓ_+ is realizable in *T*. This allows us to approximate ℓ_+ by simple closed curves σ on *S* and estimate the translation length $l_i(\sigma)$ of σ in \mathbb{H}^3/Γ_i . The estimate thus obtained contradicts a classical estimate of $l_i(\sigma)$ due to Ahlfors obtained in terms of the length of the geodesic realization of σ in τ_i^+ and τ_i^- . This final contradiction proves Theorem 14.

Finally, to hyperbolize an atoroidal 3-manifold fibering over the circle with monodromy ϕ , one picks a base Riemann surface τ , and sets $\tau_i^+ = \phi^i(\tau)$ and $\tau_i^- = \phi^{-i}(\tau)$. Then ℓ_+, ℓ_- turn out to be the stable and unstable laminations of ϕ . The 3-manifold M obtained from the double limit theorem is easily seen to be invariant under ϕ , and hence M admits a quotient which is the required hyperbolic 3-manifold.

5 Combination theorems in geometric group theory: hyperbolic groups

The fundamental combination theorem in the context of hyperbolic groups in the sense of Gromov [Gro85] is due to Bestvina and Feighn [BF92]. The theorem was motivated by Thurston's combination Theorem 9. In the context of geometric group theory, free products with amalgamation and HNN extensions can be treated on

a common footing by passing to the universal cover and looking at the resulting Bass-Serre tree of spaces [SW79]. Thus, while the main combination theorem of [BF92] provides only a weaker conclusion than Theorem 9 inasmuch as it establishes Gromov-hyperbolicity, the context is considerably more general and works for trees of spaces. It turns out that the sufficient condition in [BF92] are also necessary and this converse direction was established by Gersten [Ger98], Bowditch [Bow07] and others. The paper [BF92] spawned considerable activity in geometric group theory and have been giving rise to a number of combination theorems [Dah03, Ali05, MR08, GH09, GW11, MS12, Gau16, MO21] right up to the time of writing this article. A forthcoming book of Kapovich and Sardar [KS21] furnishes a definitive account and rather general versions of the material in Sections 5.1 and 5.2.

5.1 Trees of spaces

The framework of [BF92] is that of a tree of spaces. We follow the exposition in [Mj20] to define the relevant notions.

Definition 12 [BF92] Let (X, d) be a geodesic metric space. Let T be a simplicial tree. Let $\mathcal{V}(T)$ and $\mathcal{E}(T)$ denote the vertex set and edge set of T respectively. Then $P: X \to T$ is said to be a tree of geodesic metric spaces satisfying the *quasi-isometrically embedded condition* (or simply, the *qi condition*) if there exists a map $P: X \to T$, and constants $K \ge 1, \epsilon \ge 0$ satisfying the following:

- 1. $\forall v \in \mathcal{V}(T), X_v = P^{-1}(v) \subset X$ equipped with the induced path metric d_v is a geodesic metric space X_v . Also, the inclusion maps $i_v : X_v \to X$ are uniformly proper, i.e. $\forall M > 0, v \in T$ and $x, y \in X_v$, there exists N > 0 such that $d(i_v(x), i_v(y)) \leq M$ implies $d_v(x, y) \leq N$.
- 2. Let $e = [v_1, v_2] \in \mathcal{E}(T)$ with initial and final vertices v_1 and v_2 respectively (we assume that all edges have length 1). Let X_e be the pre-image under P of the mid-point of e. There exist continuous maps $f_e : X_e \times [v_1, v_2] \to X$, such that $f_e|_{X_e \times (v_1, v_2)}$ is an isometry onto the pre-image of the interior of e equipped with the path metric d_e .

Further, we demand that f_e is fiber-preserving, i.e. projection to the second coordinate in $X_e \times [v_1, v_2]$ corresponds via f_e to projection to the tree $P : X \to T$.

3. $f_e|_{X_e \times \{v_1\}}$ and $f_e|_{X_e \times \{v_2\}}$ are (K, ϵ) -quasi-isometric embeddings into X_{v_1} and X_{v_2} respectively. We shall often use the shorthand f_{e,v_1} and f_{e,v_2} for $f_e|_{X_e \times \{v_1\}}$ and $f_e|_{X_e \times \{v_2\}}$ respectively.

We shall refer to K, ϵ as the constants or parameters of the qi-embedding condition.

If there exists $\delta > 0$ such that the vertex and edge spaces X_v, X_e above are all δ -hyperbolic metric spaces for all vertices v and edges e of T, then $P : X \to T$ will be called *a tree of hyperbolic metric spaces*.

Definition 13 [BF92] A continuous map $f : [-k,k] \times I \to X$ is called a *hallway* of length 2k if it satisfies the following:

1. $f^{-1}(\cup X_v : v \in T) = \{-k, \cdots, k\} \times I$

2. *f* is transverse, relative to condition (1) to $\cup_e X_e$.

3. *f* maps $i \times I$ to a geodesic in X_v for some vertex space X_v .

Definition 14 [BF92] A hallway $f : [-k,k] \times I \to X$ is said to be ρ -thin if

$$d(f(i,t), f(i+1,t)) \le \rho$$

for all i, t.

A hallway $f : [-m, m] \times I \to X$ is called λ -hyperbolic if

 $\lambda l(f(\{0\} \times I)) \le \max \{ l(f(\{-m\} \times I)), l(f(\{m\} \times I)) \}.$

The girth of the hallway is defined to be the quantity

 $\min_{i} \{ l(f(\{i\} \times I)) \}.$

A hallway is *essential* if the edge path in *T* resulting from projecting the hallway under $P \circ f$ onto *T* does not backtrack (and is therefore a geodesic segment in the tree *T*).

Definition 15 (Hallways flare condition [BF92]:) The tree of spaces, *X*, is said to satisfy the *hallways flare* condition if there exist $\lambda > 1$ and $m \ge 1$ such that the following holds:

 $\forall \rho > 0$ there exists $H(=H(\rho))$ such that any ρ -thin essential hallway of length 2m and girth at least H is λ -hyperbolic.

The constants λ , *m* are referred to as the constants or parameters of the hallways flare condition. If, further, the constant ρ is fixed, then *H* will also be called a constant or parameter of the hallways flare condition.

With these notions in place, we can state the main geometric combination theorem of [BF92]:

Theorem 15 Let $P : X \to T$ be a tree of hyperbolic spaces satisfying the qiembedded condition (as in Definition 12). Further, suppose that the hallways flare condition (as in Definition 15) is satisfied. Then X is hyperbolic.

The proof of Theorem 15 in [BF92] proceeds by establishing a linear isoperimetric inequality ensuring hyperbolicity. We shall indicate a different proof scheme below in the special case that the edge-to-vertex inclusion maps are uniform quasi-isometries rather than qi-embeddings. The forthcoming book [KS21] provides a new proof as well.

16

5.2 Metric Bundles

The notion of a metric bundle [MS12] adapts the idea of a fiber bundle to a coarse geometric context. We shall describe below the main combination theorem of [MS12] which is an analog of Theorem 15 in this context.

Definition 16 Let (X, d_X) and (\mathcal{B}, d_B) be geodesic metric spaces. Let $c, K \ge 1$ be constants and $h : \mathbb{R}^+ \to \mathbb{R}^+$ a function. $P : X \to \mathcal{B}$ is called an (h, c, K)-*metric bundle* if

- 1. *P* is 1-Lipschitz.
- 2. For each $z \in \mathcal{B}$, $F_z = P^{-1}(z)$ is a geodesic metric space with respect to the path metric d_z induced from (X, d_X) . We refer to F_z as the *fiber* over z. We further demand that the inclusion maps $i_z : (F_z, d_z) \to X$ are uniformly metrically proper as measured with respect to h, i.e. for all $z \in \mathcal{B}$ and $u, v \in F_z$,

 $d_X(i_z(u), i_z(v)) \le N$ implies that $d_z(u, v) \le f(N)$.

- 3. For $z_1, z_2 \in \mathcal{B}$ with $d_B(z_1, z_2) \leq 1$, let γ be a geodesic in \mathcal{B} joining them. Then for any $z \in \gamma$ and $x \in F_z$, there is a path in $p^{-1}(\gamma)$ of length at most *c* joining *x* to both F_{z_1} and F_{z_2} .
- 4. For $z_1, z_2 \in \mathcal{B}$ with $d_B(z_1, z_2) \leq 1$ and $\gamma \subset \mathcal{B}$ a geodesic joining them, let $\phi : X_{z_1} \to X_{z_2}$, be any map such that for all $x_1 \in X_{z_1}$ there is a path of length at most *c* in $P^{-1}(\gamma)$ joining x_1 to $\phi(x_1)$. Then ϕ is a *K*-quasi-isometry.

If in addition, there exists δ' such that each X_z is δ' -hyperbolic, then $P : X \to \mathcal{B}$ is called an (h, c, K)- metric bundle of δ' -hyperbolic spaces (or simply a metric bundle of hyperbolic spaces if the constants are implicit).

It is pointed out in [MS12] that condition (4) follows from the previous three (with suitable K); but it is more convenient to have it as part of our definition.

A closely related notion of a *metric graph bundle* often turns out to be more useful:

Definition 17 [MS12, Definition 1.2] Suppose X and \mathcal{B} are metric graphs and $f : \mathbb{N} \to \mathbb{N}$ is a proper function. We say that X is an *f*-metric graph bundle over \mathcal{B} if there exists a surjective simplicial map $\pi : X \to \mathcal{B}$ such that the following hold.

- 1. For all $b \in V(\mathcal{B})$, $\mathcal{F}_b := \pi^{-1}(b)$ is a connected subgraph of X. Moreover, the inclusion maps $\mathcal{F}_b \to X$, $b \in V(\mathcal{B})$ are uniformly metrically proper as measured by f.
- 2. For all adjacent vertices $b_1, b_2 \in V(\mathcal{B})$, any $x_1 \in V(\mathcal{F}_{b_1})$ is connected by an edge to some $x_2 \in V(\mathcal{F}_{b_2})$.

For all $b \in V(\mathcal{B})$, F_b is called the *fiber* over *b* and its path metric is denoted by d_b . It is pointed out in [MS12] that any metric bundle is quasi-isometric to a metric graph bundle, where the quasi-isometry coarsely preserves fibers and restricts to a quasi-isometry of fibers. Condition (2) of Definition 17 immediately shows that if $\pi : X \to \mathcal{B}$ is a metric graph bundle then for any points $v, w \in V(\mathcal{B})$ we have $Hd(F_v, F_w) < \infty$.

Example Let

$$1 \to N \to G \to Q \to 1$$

be an exact sequence of finitely generated groups. Choose a finite generating set of N and extend it to a finite generating set of G. The image of the finite generating set of G in Q under the quotient map is then a generating set of Q. This gives a natural simplicial map $P : \Gamma_G \to \Gamma_Q$ between the respective Cayley graphs. This is the prototypical example of a metric graph bundle. The fibers are all copies of Γ_N . \Box

Definition 18 Suppose X is an *f*-metric graph bundle over \mathcal{B} . Given $k \ge 1$ and a connected subgraph $\mathcal{A} \subset \mathcal{B}$, a *k*-qi section over \mathcal{A} is a map $s : \mathcal{A} \to X$ such that s is a *k*-qi embedding and $\pi \circ s$ is the identity map on \mathcal{A} .

For any hyperbolic metric space F with more than two points in its Gromov boundary ∂F , there is a coarsely well-defined *barycenter map*

$$\phi:\partial^3 F\to F$$

mapping an unordered triple (a, b, c) of distinct points in ∂F to a centroid of the ideal triangle spanned by (a, b, c). We shall say that the barycenter map $\phi : \partial^3 F \to F$ is *N*-coarsely surjective if *F* is contained in the *N*-neighborhood of the image of ϕ . A *K*-*qi*-section $\sigma : \mathcal{B} \to X$ is a *K*-*qi*-embedding from \mathcal{B} to X such that $P \circ \sigma$ is the identity map. The following guarantees the existence of qi-sections for metric bundles:

Proposition [MS12, Section 2.1] Given δ , N, c, $K \ge 0$ and proper $f : \mathbb{N} \to \mathbb{N}$, there exists K_0 such that the following holds.

Let $P : X \to \mathcal{B}$ be an (f, c, K)-metric bundle of δ -hyperbolic spaces such that all barycenter maps $\phi_b : \partial^3 F_b \to F_b$ are *N*-coarsely surjective, Then through each point of *X*, there exists a K_0 -qi section.

A similar statement holds for metric graph bundles.

The following gives the analog of Definition 15 in the context of metric bundles and metric graph bundles:

Definition 19 Let $P : X \to \mathcal{B}$ be a metric bundle or a metric graph bundle. $P : X \to \mathcal{B}$ is said to satisfy a *flaring condition* if $\forall k \ge 1$, there exist $\lambda_k > 1$ and $n_k, M_k \in \mathbb{N}$ such that the following holds:

Let $\gamma : [-n_k, n_k] \to \mathcal{B}$ be a geodesic and let $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ be two *k*-qi sections of γ in *X*. If $d_{\gamma(0)}(\tilde{\gamma}_1(0), \tilde{\gamma}_2(0)) \ge M_k$, then

 $\lambda_k.d_{\gamma(0)}(\tilde{\gamma_1}(0),\tilde{\gamma_2}(0)) \le \max\{d_{\gamma(n_k)}(\tilde{\gamma_1}(n_k),\tilde{\gamma_2}(n_k)), d_{\gamma(-n_k)}(\tilde{\gamma_1}(-n_k),\tilde{\gamma_2}(-n_k))\}.$

The following Theorem is the analog of Theorem 15 in the context of metric (graph) bundles.

Theorem 18 Suppose that $P : X \to \mathcal{B}$ is a metric bundle or a metric graph bundle such that all fibers F_z are uniformly hyperbolic, and the barycenter maps are uniformly coarsely surjective. Equivalently, by Proposition 17, there exists $\rho \ge 1$ such that for every $x \in X$, there exists a ρ -qi section $s : \mathcal{B} \to X$ passing through x, i.e. $s \circ P(x) = x$.

Then if X satisfies the qi-embedded condition and the flaring condition (as in Definition 19) corresponding to ρ -qi sections, then X is hyperbolic.

Conversely, if X is hyperbolic, then as a metric bundle or metric graph bundle, X satisfies the flaring condition.

5.2.1 Ladders

A tool that has turned out to be considerably useful in the context of both trees of spaces and metric bundles is the notion of a ladder. In particular, for our proof of Theorem 18 (sketched in Section 5.2.2), we shall use it. The notion is related to, but different from that of a hallway. Ladders were introduced in [Mit98b] in the context of trees of spaces and in [Mit98a] in the context of groups. Instead of going through the construction in detail, we extract the relevant features from the ladder construction of [Mit98b, Mit98a]. The following is a restatement of [Mit98b, Theorem 3.6] reformulated to emphasize the connection with hallways.

Theorem 19 Given $\delta \ge 0, K \ge 1, \epsilon \ge 0$ there exists *D* such that the following holds. We consider one of the two following situations:

- 1. $P: X \to T$ is a tree of δ -hyperbolic spaces as in Definition 12 with parameters K, ϵ . Let F_v be a vertex space,
- 2. $P: X \to \mathcal{B}$ is a metric bundle or metric graph bundle, and F_v is a fiber.

In both cases, the intrinsic metric on F_v is denoted by d_v . Then for every geodesic segment $\mu \subset (F_v, d_v)$ there exists a D-qi-embedded subset \mathcal{L}_{μ} of X such that the following holds.

- 1. $F_v \cap \mathcal{L}_\mu = \mu$,
- 2. (a) For $P: X \to T$ a tree of hyperbolic metric spaces and every $w \in T$, $F_w \cap \mathcal{L}_\mu$ is either empty or a geodesic μ_w in (F_w, d_w) . Further, there exists a subtree $\mathcal{T}_1 \subset T$ such that the collection of vertices $w \in T$ satisfying $F_w \cap \mathcal{L}_\mu \neq \emptyset$ equals the vertex set of T_1 .

(b) For $P: X \to \mathcal{B}$ a metric bundle or metric graph bundle, $F_w \cap \mathcal{L}_\mu$ is a geodesic μ_w in (F_w, d_w) .

3. There exists $\rho_0 \ge 1$ such that through every $z \in \mathcal{L}_{\mu}$, there exists a ρ_0 -qi-section σ_z of [v, P(z)] contained in \mathcal{L}_{μ} satisfying

$$\sigma_z(P(z)) = z, \quad \sigma_z(v) \in \mu.$$

Further, there exists a D-coarse Lipschitz retraction $\Pi_{\mu}: X \to \mathcal{L}_{\mu}$, i.e.

1.
$$d(\Pi_{\mu}(x), \Pi_{\mu}(y)) \le Dd(x, y) + D, \forall x, y \in X,$$

2. $\Pi_{\mu}(x) = x, \forall x \in \mathcal{L}_{\mu}.$

The qi-embedded set \mathcal{L}_{μ} is called a *ladder* in [Mit98a, Mit98b]. Theorem 19 shows in particular that there is a (2D, 2D)- quasigeodesic of (X, d_X) joining the end-points of μ and lying on \mathcal{L}_{μ} .

Remark Note that in Theorem 19, we have *not* assumed that X is hyperbolic: no assumptions on the global geometry of X are necessary here. \Box

5.2.2 Idea behind the proof of Theorem 18

We focus on the metric graph bundle case for convenience. Theorem 19 guarantees that for any pair of points x, y in a metric graph bundle X, there exist

- 1. Qi-sections Σ_x, Σ_y through x, y.
- 2. A ladder $\mathcal{L}(x, y)$ bounded by Σ_x, Σ_y . In fact, in this case (as shown in [Mit98a, MS12]), $\mathcal{L}(x, y) \cap F_b$ equals a geodesic in F_b joining $\Sigma_x(b), \Sigma_y(b)$ (here we are abusing notation slightly by identifying the qi-sections Σ_x, Σ_y with their images).

Thus, for every $x, y \in X$ there are preferred quasigeodesics in X contained in $\mathcal{L}(x, y)$. We have not used the flaring condition so far. The flaring condition guarantees hyperbolicity of $\mathcal{L}(x, y)$. We shall return to this shortly. Hyperbolicity of $\mathcal{L}(x, y)$ ensures (by the Morse Lemma) that all quasigeodesics in $\mathcal{L}(x, y)$ joining x, y are in a bounded neighborhood of each other. This gives a family of paths in X, one for every pair x, y. We then use a path-families argument following Hamenstadt [Ham05] and a criterion due to Bowditch to conclude that X is hyperbolic.

We return to the proof of hyperbolicity of $\mathcal{L}(x, y)$. We note that $\mathcal{L}(x, y)$ is a bundle over \mathcal{B} where the fibers are intervals. The flaring condition is inherited by $\mathcal{L}(x, y)$ with slightly worse constants. Thus, we are reduced to proving Theorem 18 in the special case that fibers are intervals. To do this, we decompose the ladder $\mathcal{L}(x, y)$ using qi-sections contained in $\mathcal{L}(x, y)$ into a finite number of ladders 'stacked one on top of another'. Thus, there exist disjoint sections $\Sigma_x = \Sigma_0, \Sigma_1, \dots, \Sigma_n = \Sigma_y$ and ladders \mathcal{L}_i bounded by Σ_{i-1}, Σ_i such that distinct \mathcal{L}_i 's have disjoint interiors. The ubiquity of qi-sections allows us to ensure that each of these smaller ladders has bounded girth (in the spirit of Definition 14), i.e. Σ_{i-1}, Σ_i are at a bounded distance from each other along some fiber F_b and flare away from each other as one goes to infinity in *B*. A further path families argument following [Ham05] allows us to prove that $\mathcal{L}(x, y)$ is hyperbolic.

A word about the proof sketch above. Note that we use only the 1-dimensional property of quasigeodesics flaring and path families to prove the combination theorem in this case, as opposed to the more '2-dimensional' area argument of [BF92]. This has been considerably refined in [KS21] to give a new path-families proof of Theorem 15.

5.3 Relatively hyperbolic combination theorems

We refer the reader to [Far98, Gro85, Bow12] for the basics of relative hyperbolicity. Theorem 15 was generalized to the context of trees of relatively hyperbolic spaces in two different ways:

- 1. Using an acylindricity hypothesis in [Dah03] and [Ali05]. This is in the spirit of Theorem 10.
- 2. Using the flaring condition in [MR08, Gau16]. This in the spirit of Theorem 14.

5.3.1 Relatively hyperbolic combination theorem using acylindricity

Let *G* be hyperbolic relative to a finite collection $\mathcal{P} = \{P_1, \dots, P_k\}$ of parabolic subgroups. Let $\partial_h G$ denote the Bowditch boundary of *G*. Let $H \subset G$ be a relatively quasiconvex subgroup [Hru10]. We shall give Dahmani's version of the combination theorem [Dah03] below. Let $\Lambda_H \subset \partial_h G$ denote the limit set of *H*. A relatively quasiconvex subgroups *H* is *full relatively quasi-convex* if it is quasi-convex and if, for any infinite sequence $g_n \in G$ in distinct left cosets of *H*, the intersection $\cap_n g_n(\Lambda_H)$ is empty.

Lemma 1 [Dah03, Lemma 1.7] Let G be hyperbolic relative to a finite collection $\mathcal{P} = \{P_1, \dots, P_k\}$ of parabolic subgroups. Let H be a full relatively quasi-convex subgroup. Let P be a conjugate of one of the P_i 's. Then $P \cap H$ is either finite, or of finite index in P.

Definition 20 [Sel97] The action of a group *G* on a tree *T* is *k*-acylindrical for some $k \in \mathbb{N}$ if the stabilizer of any geodesic of length *k* in *T* is finite. The action of a group *G* on a tree *T* is acylindrical if it is *k*-acylindrical for some $k \in \mathbb{N}$

A finite graph of groups is said to be acylindrical, if the action on the associated Bass-Serre tree is acylindrical.

Then Dahmani's combination theorem states:

Theorem 21 [Dah03] Let G be the fundamental group of an acylindrical finite graph of relatively hyperbolic groups, whose edge groups are full quasi-convex subgroups of the adjacent vertex groups. Let G be the family of images of the maximal parabolic subgroups of the vertex groups, and their conjugates in G. Then G is strongly hyperbolic relative to G.

The approach in [Dah03] is quite different from that of [BF92]. From the Bowditch boundaries of the vertex and edge groups, a metrizable compact space Z is constructed in such a way that G naturally acts on Z. It is then shown that this action is a convergence action. Finally, it is shown that the action is geometrically finite, forcing G to have a relatively hyperbolic structure.

The Bass-Serre tree of G has vertex groups G_v and edge groups G_e . Hence, associated to the Bass-Serre tree T there is a natural tree (T) of compact spaces

given by $\partial_h G_v$ and $\partial_h G_e$. The set *Z* is built [Dah03, Section 2] from these copies of $\partial_h G_v$ and $\partial_h G_e$. Suppose $e = [v_1, v_2]$ is an edge of *T*. For all such edges *e*, glue together $\partial_h G_{v_1}$ and $\partial_h G_{v_2}$ along the limit set $\partial_h G_e$. The relevant identification space is thus obtained from the set $\sqcup_{v \in V(T)} \partial_h G_v$ by identifying pairs of points according to the images of $\partial_h G_e$. Finally, the base tree *T* encodes (infinite) directions that are 'transverse' to all the vertex spaces. The set *Z* is then obtained from topologizing $\partial T \cup \sqcup_{v \in V(T)} \partial_h G_v / \sim$, where ~ is the equivalence relation given by edge spaces.

Alibegovic [Ali05] proves a similar combination theorem for relatively hyperbolic groups following the original strategy of Bestvina and Feighn in Theorem 15 using the linear isoperimetric inequality characterization of hyperbolicity.

5.3.2 Relatively hyperbolic combination theorem using flaring

We next define a tree of relatively hyperbolic spaces in general.

Definition 21 [MR08] A tree $P : X \to \mathcal{T}$ of geodesic metric spaces is said to be a tree of relatively hyperbolic metric spaces if in addition to the conditions of Definition 12

- (4) each vertex space X_v is strongly hyperbolic relative to a collection of subsets \mathcal{H}_v and each edge space X_e is strongly hyperbolic relative to a collection of subsets \mathcal{H}_e . The sets $H_{v,\alpha} \in \mathcal{H}_v$ or $H_{e,\alpha} \in \mathcal{H}_e$ are referred to as *horosphere-like sets*.
- (5) the maps f_{e,v_i} above, for i = 1, 2, are *strictly type-preserving*. That is, for i = 1, 2and for any $H_{v_i,\alpha} \in \mathcal{H}_{v_i}, f_{e,v_i}^{-1}(H_{v_i,\alpha})$, is either empty or equals some $H_{e,\beta} \in \mathcal{H}_e$. Further, for all $H_{e,\beta} \in \mathcal{H}_e$, there exists v and $H_{v,\alpha}$, such that $f_{e,v}(H_{e,\beta}) \subset H_{v,\alpha}$.
- (6) There exists $\delta > 0$ such that each $\mathcal{E}(X_{\nu}, \mathcal{H}_{\nu})$ is δ -hyperbolic (here, $\mathcal{E}(X_{\nu}, \mathcal{H}_{\nu})$ denotes the electric space obtained from X_{ν} by electrifying all the horospherelike sets in \mathcal{H}_{ν}).
- (7) The induced maps of the coned-off edge spaces into the coned-off vertex spaces $\widehat{f_{e,v_i}} : \mathcal{E}(X_e, \mathcal{H}_e) \to \mathcal{E}(X_{v_i}, \mathcal{H}_{v_i}) \ (i = 1, 2)$ are uniform quasi-isometries. This is called the *qi-preserving electrification condition*

We state conditions (4) and (6) in conjunction by saying that X_v is *strongly* δ -hyperbolic relative to \mathcal{H}_v .

We explain condition (7) briefly. Given the tree of spaces $P : X \to T$ with vertex spaces X_v and edge spaces X_e there exists a naturally associated tree whose vertex spaces are the electrified spaces $\mathcal{E}(X_v, \mathcal{H}_v)$ and edge spaces are the electrified spaces $\mathcal{E}(X_e, \mathcal{H}_e)$ obtained by electrifying the respective horosphere like sets. Condition (4) of the above definition ensures that we have natural inclusion maps of edge spaces $\mathcal{E}(X_e, \mathcal{H}_e)$ into adjacent vertex spaces $\mathcal{E}(X_v, \mathcal{H}_v)$. The resulting tree of coned-off spaces $P : \mathcal{T}C(X) \to \mathcal{T}$ is referred to simply as the *induced tree of coned-off spaces*. The *cone locus* of $\mathcal{T}C(X)$ is the forest given by the following:

1. the vertex set $\mathcal{V}(\mathcal{T}C(X))$ consists of the cone-points $\{c_{\nu,\alpha} \text{ in the vertex spaces } X_{\nu} \text{ resulting from the electrification operation of the horosphere-like sets } H_{\nu,\alpha} \in \mathcal{H}_{\nu}.$

2. the edge set $\mathcal{E}(\mathcal{TC}(X))$ consists of the cone-points $\{c_{e,\alpha} \text{ in the edge set } X_e \text{ resulting from the electrification operation of the horosphere-like sets <math>H_{e,\alpha} \in \mathcal{H}_e$..

Each connected component of the cone-locus is a maximal cone-subtree. The collection of maximal cone-subtrees is denoted by $C\mathcal{T}$ and elements of $C\mathcal{T}$ are denoted as CT_{α} . Note that each maximal cone-subtree CT_{α} naturally gives rise to a tree CT_{α} of horosphere-like subsets depending on which cone-points arise as vertices and edges of CT_{α} . The metric space that CT_{α} gives rise to is denoted as C_{α} . We refer to any such C_{α} as a maximal cone-subtree of horosphere-like spaces. The induced tree of horosphere-like sets is denoted by

$$g_{\alpha}: C_{\alpha} \to CT_{\alpha}.$$

The collection of these maps will be denoted as \mathcal{G} . The collection of the maximal cone-subtree of horosphere-like spaces C_{α} is denoted as C. Note thus that each CT_{α} thus appears both as a subset of $\mathcal{TC}(X)$ as well as the underlying tree of C_{α} .

Definition 22 (Cone-bounded hallways strictly flare condition:)

An essential hallway of length 2*k* is *cone-bounded* if $f(i \times \partial I)$ lies in the conelocus for $i = \{-k, \dots, k\}$.

The tree of spaces, X, is said to satisfy the *cone-bounded hallways flare* condition if there are numbers $\lambda > 1$ and $k \ge 1$ such that any cone-bounded hallway of length 2k is λ -hyperbolic, where λ, k are called the constants or parameters of the strict flare condition.

We now state the combination theorem for relative hyperbolicity using the flaring condition.

Theorem 22 [MR08, Gau16] Let $P : X \to T$ be a tree of uniformly relatively hyperbolic spaces in the sense of Definition 21 satisfying the qi-embedded condition, such that the resulting tree of coned-off spaces satisfies

1. the hallways flare condition,

2. the cone-bounded hallways flare condition.

Then X is hyperbolic relative to the maximal cone-subtrees of horosphere-like spaces.

5.4 Effective quasiconvexity and flaring

We now state a couple of theorems along the lines of Theorem 15 and 18 ensuring quasiconvexity of a subspace of a vertex space. The first, due to Ilya Kapovich [Kap01a] is in the setup of an acylindrical graph of groups:

Theorem 23 Let G be a finite acylindrical graph of groups where all vertex and edge groups are hyperbolic and edge-to-vertex inclusions are quasi-isometric embeddings. Let $T \subset G$ be a maximal subtree. Let G denote the group corresponding to the tree T. (By Theorem 15, G is hyperbolic.) Then each vertex group G_v of G is quasiconvex in G.

Next, we shall consider in a unified way the two following situations:

- P: X → T is a tree of hyperbolic metric spaces satisfying the qi-embedded condition with constants K, ε and the hallways flare condition with parameters λ₀, m₀. Further, if ρ₀ is given we shall assume an additional constant H₀ as a lower bound for girths of ρ₀-thin hallways.
- 2. $P : X \to \mathcal{B}$ is a metric bundle or a metric graph bundle satisfying the flaring condition with constants as i Definitions 16, 17, 19.

Also (X_v, d_v) will, be a vertex space of X (in the tree of spaces case) or $P^{-1}(v)$ equipped with the induced metric in the metric (graph) bundle case.

Definition 23 Let $P : X \to \mathcal{T}$ be a tree of hyperbolic spaces. Let $Y \subset (X_v, d_v)$ be a *C*-quasiconvex subset of (X_v, d_v) . We say that *Y* flares in all directions with *parameter K* if for any geodesic segment $[a, b] \subset (X_v, d_v)$ with $a, b \in Y$ and any ρ -thin hallway $f : [0, k] \times I \to X$ satisfying

1.
$$\rho \le \rho_0$$
,
2. $f(\{0\} \times I) = [a, b]$,
3. $l([a, b]) \ge K$,
4. $k \ge K$,

the length of $f(\{k\} \times I)$ satisfies

$$l(f(\{k\} \times I)) \ge \lambda l([a, b]).$$

Similarly, let $P : X \to \mathcal{B}$ be a metric bundle or metric graph bundle with hyperbolic fiber. Let $Y \subset X_v$ be quasiconvex. Further, assume that there is a ρ -qi section through every $x \in X$ (cf. the second hypothesis of Theorem 18).

We say that *Y* flares in all directions with parameter $K \ge 0, D \ge 1, \lambda > 1$ if the following holds:

Let $\gamma : [0, D] \to \mathcal{B}$ be a geodesic such that $\gamma(0) = v$ and let $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ be two ρ -qi lifts (sections) of γ in X. If $d_v(\tilde{\gamma}_1(0), \tilde{\gamma}_2(0)) \ge K$, then we have

$$\lambda.d_{\nu}(\tilde{\gamma}_1(0),\tilde{\gamma}_2(0)) \le d_{\gamma(D)}(\tilde{\gamma}_1(D),\tilde{\gamma}_2(D)).$$

We can now state the a Proposition guaranteeing quasiconvexity of subsets of vertex spaces.

Proposition [Mj20] Given K, C, there exists C_0 such that the following holds. Let $P : X \to T$ and X_v be as in Theorem 19 above. If Y is a C-quasiconvex subset of (X_v, d_v) and flares in all directions with parameter K, then Y is C_0 -quasiconvex in (X, d_X) .

Conversely, given C_0 , there exist K, C such that the following holds. For $P : X \to \mathcal{T}$ and X_v as above, if $Y \subset X_v$ is C_0 -quasiconvex in (X, d_X) , then it is a *C*-quasiconvex subset in (X_v, d_v) and flares in all directions with parameter K. \Box

A similar statement holds for metric (graph) bundles.

24

Proposition Given K, D, λ, C , there exists C_0 such that the following holds.

Let $P : X \to \mathcal{B}$ be a metric (graph) bundle and X_v be as in Definition 23. If Y is a *C*-quasiconvex subset of (X_v, d_v) and flares in all directions with parameters K, D, λ , then Y is C_0 -quasiconvex in (X, d_X) .

Conversely, given C_0 , there exist K, D, λ, C such that the following holds. For $P : X \to \mathcal{B}$ a metric (graph) bundle and X_v as above, if $Y \subset X_v$ is C_0 -quasiconvex in (X, d_X) , then it is a *C*-quasiconvex subset in (X_v, d_v) and flares in all directions with parameters K, D, λ .

6 Combination theorems in geometric group theory: cubulations

We turn now to the remarkable work during the last decade on special cube complexes. We refer to [HW08] for the basics of special cube complexes. Let \mathcal{G} denote a finite graph, $RAAG(\mathcal{G})$ the right-angled Artin group associated to \mathcal{G} , and $\mathcal{S}(\mathcal{G})$ its Salvetti complex. A cube complex C is said to be *special* if there exists a combinatorial local isometry from C to $\mathcal{S}(\mathcal{G})$ for some finite graph \mathcal{G} . By Agol's resolution of Wise's conjecture in [Ago13] (see Theorem 30 below), hyperbolic groups that are virtually special are precisely those that act geometrically on a CAT(0) cube complex. We give a brief account of some of the combination theorems that have been proved around this theme.

In [HW10], Hsu and Wise proved the precursor of all virtually special combination theorems by showing that if a hyperbolic group G splits as a finite graph of finitely generated free groups with cyclic edge groups, then G is virtually special. In [HW15b], they later generalized this to amalgamated products of free groups over a finitely generated malnormal subgroup. A landmark combination theorem due to Haglund and Wise concerns the combination of hyperbolic virtually special cubulable groups along malnormal quasiconvex subgroups:

Theorem 26 [HW12] Let A, B, M be compact virtually special cube complexes. Suppose that $G_A = \pi_1(A)$, $G_B = \pi_1(B)$, and $G_M = \pi_1(M)$ are hyperbolic. Let $M \xrightarrow{i_A} A$, and $M \xrightarrow{i_B} B$ be local isometries of cube complexes such that $i_{A*}(G_M)$ and $i_{B*}(G_M)$ are quasiconvex and malnormal in G_A and G_B respectively. Let $X = A \cup_M B$ be the cube complex obtained by gluing A and B together along M using $M \times [0, 1]$. Then X is virtually special.

Theorem 26 generalizes earlier work of Wise [Wis02] where he showed that any 2-complex built by amalgamating (in terms of fundamental group) two finite graphs along a malnormal immersed graph is virtually special. Theorem 26 is also a crucial ingredient in Wise's proof of the virtual specialness of hyperbolic groups admitting a quasiconvex hierarchy. This is very much in the spirit of the Haken hierarchy for Haken 3-manifolds and Thurston's hyperbolization of such manifolds (cf. Section 4.1).

Theorem 27 [Wis21] Let G be a hyperbolic group admitting a quasiconvex hierarchy. Then G is the fundamental group of a compact non-positively curved cube complex that is virtually special.

We list below some of the important consequences of Theorem 27. The following resolved a conjecture of Baumslag:

Theorem 28 [Wis21] Every one-relator group with torsion is virtually special.

In the context of hyperbolic 3-manifolds, Wise showed the following.

Theorem 29 [Wis21] Compact hyperbolic Haken manifolds are virtually special.

Using work of Kahn and Markovic [KM12], Bergeron and Wise [BW12] proved that all hyperbolic 3-manifolds can be cubulated, i.e. they act geometrically on CAT(0) cube complexes. This led Wise to conjecture that hyperbolic groups that act geometrically on CAT(0) cube complexes are virtually special. The following celebrated theorem of Agol resolved this conjecture affirmatively:

Theorem 30 [Ago13] Hyperbolic groups acting geometrically on CAT(0) cube complexes are virtually special.

A flurry of activity ensued in trying to show that several naturally defined hyperbolic groups are, in fact, cubulable. In [HW16, HW15a], Hagen and Wise proved that hyperbolic groups G admitting an exact sequence of the form

$$1 \to F_n \to G \to \mathbb{Z} \to 1$$

are cubulable. (Here F_n denotes the free group on *n* generators.) Hence, by Agol's Theorem 30, such groups *G* are virtually special. In a different direction, Manning, the first author and Sageev [MMS19] showed that there exist cubulable hyperbolic groups *G* admitting an exact sequence of the form

$$1 \to \pi_1(S) \to G \to F_n \to 1$$
,

where S is a closed surface of genus greater than one. Again, by Agol's Theorem 30, such groups G are virtually special.

Finally, we mention work of Przytycki and Wise [PW18], who proved the virtual specialness of fundamental groups of 3-manifolds whose JSJ decomposition has both a hyperbolic as well as a Seifert-fibered piece. 3-manifolds admitting such a JSJ decomposition are called *mixed*. As a consequence of their result, the authors show that mixed manifolds virtually fiber.

The proof in [PW18] proceeds by first showing that there are enough codimension one surface subgroups to ensure cubulability. This is established by combining surfaces coming from the graph manifold pieces with those coming from hyperbolic pieces. Once cubulability has been established, the malnormal special quotient theorem [Wis21] (see also [AGM16]) is used to establish specialness of the cube complex thus built. In this section, we have given only a cursory treatment of a topic that, starting with [Sag95] has undergone tremendous development over the last two decades. We refer the reader to the books [AFW15] and [Wis12] for a more comprehensive treatment.

7 Holomorphic dynamics and polynomial mating

Before entering the theme of combination theorems in holomorphic dynamics, we say a few words on the history of the subject, and sketch briefly some of the philosophical parallels between holomorphic dynamics and Kleinian groups and some of the developments inspired by this synergy. These will also serve as a motivation for combination theorems involving complex polynomials and Kleinian groups that we will discuss later in the section.

7.1 Historical comments

The study of dynamics of rational maps on the Riemann sphere started with groundbreaking work of Fatou and Julia [Fat19, Fat20a, Fat20b, Fat26, Jul18, Jul22] in the 1920s. The subject remained dormant for several decades barring a handful of important contributions, most notably by Siegel [Sie42] and Brolin [Bro65]. Around the 1970s, the availability of computers allowed Feigenbaum and Mandelbrot to perform numerical experiments on finer structures of dynamical and parameter planes of real/complex-analytic maps. Their pioneering discoveries infused fresh blood into the field, and gave rise to problems and conjectures that played pivotal roles in the development of the modern theory of holomorphic dynamics.

A revolutionary contribution came from Sullivan, who introduced quasiconformal methods into the study of rational dynamics to prove nonexistence of wandering domains in the Fatou set for rational maps [Sul85]. The seminal work of Douady and Hubbard on the dynamics of quadratic polynomials and the structure of the Mandelbrot set [DH85, DH84] turned out to be equally fundamental in that the techniques devised by them were robust enough to be applied to the study of a wide variety of holomorphic dynamical systems.

Sullivan proposed a dictionary between Kleinian groups and rational dynamics that was motivated by various common features shared by them [Sul85, p. 405]. In addition to the apparent similarities between the topological structures of the *limit set* (respectively, the *domain of discontinuity*) of a Kleinian group and the *Julia set* (respectively, the *Fatou set*) of a rational map, there are deeper similarities between the techniques employed in proving various statements in these two parallel worlds. In fact, in the same paper, Sullivan gave a new proof of Ahlfors' finiteness theorem which closely parallels the proof of the 'no wandering Fatou component' theorem for rational maps.

Around the same time, Thurston proved a topological characterization for an important class of rational maps [DH93]. This result, which is a philosophical analog of the hyperbolization of atoroidal Haken 3-manifolds, has given rise to a wealth of rich and beautiful results that we will not be touching upon in this survey (see [Ree16, §9] and the references therein).

We should emphasize that the aforementioned dictionary is not an automatic method for translating results in one setting to the other, but rather an inspiration for results and proof techniques. We now list a few prominent pieces of work motivated by this dictionary. In [LM97], Lyubich and Minsky constructed "an explicit object that plays for a rational map the role played by the hyperbolic 3-orbifold quotient of a Kleinian group". Bullett and Penrose constructed matings of holomorphic quadratic polynomials and the modular group as *holomorphic correspondences* [BP94]. Sullivan and McMullen introduced Teichmüller spaces of conformal dynamical systems in the spirit of Teichmüller spaces of Riemann surfaces in [SM98]. McMullen established conceptual connections between renormalization ideas used in holomorphic dynamics and the study of 3-manifolds fibering over the circle [McM98]. Pilgrim proved a canonical decomposition theorem for Thurston maps as an analog of the torus decomposition theorem for 3-manifolds [Pil03]. Another noteworthy development in the framework of the above dictionary is the recent work of Luo [Luo19, Luo21a, Luo21b], where results in rational dynamics were proved using techniques that are closely related to Thurston's work on 3-manifolds.

We refer the reader to [Mil06, CG93] for a basic introduction to rational dynamics, to [Lyu17] for a comprehensive account on the dynamics of quadratic polynomials and the Mandelbrot set, to the recent survey article by Rees on major advances in the field [Ree16], and a survey by DeZotti on connections between holomorphic dynamics and other branches of mathematics [DeZ20].

7.2 Mating of polynomials

The operation of polynomial mating, which was introduced by Douady in [Dou83], constructs a rational map on $\widehat{\mathbb{C}}$ by combining the actions of two complex polynomials. Since the first appearance of the notion, several closely related definitions and perspectives have been put forward. In this survey, we will follow the route adopted in [PM12] (see [Roe12] for the original formulation and some historical comments, and [Mil04] for a lucid account of the mating construction along with a detailed worked out example).

To define the operation of polynomial mating formally, we need to introduce some terminology. The *Fatou set* of a rational map *R*, denoted by $\mathcal{F}(R)$, is the largest open subset of $\widehat{\mathbb{C}}$ on which the sequence of iterates $\{R^{\circ n}\}_{n\geq 0}$ forms a normal family. Its complement is called the *Julia set*, and is denoted by $\mathcal{J}(R)$. For a complex polynomial *P*, the *filled Julia set* (i.e., the set of points with bounded forward orbits) and the *basin of attraction of infinity* (i.e., the complement of the filled Julia set) are denoted by $\mathcal{K}(P)$ and $\mathcal{B}_{\infty}(P)$, respectively. We refer the reader to [Mil06] for basic topological and dynamical properties of these sets.

A rational map R is called *postcritically finite* if each of its critical points has a finite forward orbit. R is called *hyperbolic* if each of its critical points converges to an attracting cycle under forward iteration.

If *P* is a monic, centered polynomial of degree *d* with a connected Julia set, then there exists a conformal map $\phi_P : \widehat{\mathbb{C}} \setminus \overline{\mathbb{D}} \to \mathcal{B}_{\infty}(P)$ that conjugates z^d to *P*, and satisfies $\phi'_P(\infty) = 1$ [Mil06, Theorem 9.1, Theorem 9.5]. We will call ϕ_P the

Böttcher coordinate for *P*. Furthermore, if $\partial \mathcal{K}(P) = \mathcal{J}(P)$ is locally connected, then ϕ_P extends to a semiconjugacy between $z^d|_{\mathbb{S}^1}$ and $P|_{\mathcal{J}(P)}$. In this case, the map $\phi_P : \mathbb{S}^1 \to \mathcal{J}(P)$ is called the *Carathéodory loop/semi-conjugacy* for $\mathcal{J}(P)$.

Now let P_1, P_2 be two monic polynomials of the same degree $d \ge 2$ with connected and locally connected filled Julia sets. We consider the disjoint union $\mathcal{K}(P_1) \sqcup \mathcal{K}(P_2)$ and the map

$$P_1 \sqcup P_2 : \mathcal{K}(P_1) \sqcup \mathcal{K}(P_2) \to \mathcal{K}(P_1) \sqcup \mathcal{K}(P_2),$$

$$P_1 \sqcup P_2|_{\mathcal{K}(P_1)} = P_1, \quad P_1 \sqcup P_2|_{\mathcal{K}(P_2)} = P_2.$$

Let ~ be the equivalence relation on $\mathcal{K}(P_1) \sqcup \mathcal{K}(P_2)$ generated by $\phi_{P_1}(z) \sim \phi_{P_2}(\overline{z})$, for all $z \in \mathbb{S}^1$. It is easy to check that ~ is $P_1 \sqcup P_2$ -invariant, and hence it descends to a continuous map $P_1 \perp P_2$ to the quotient $\mathcal{K}(P_1) \perp \mathcal{K}(P_2) := (\mathcal{K}(P_1) \sqcup \mathcal{K}(P_2)) / \sim$ (see [PM12, §4.1] for details). The map $P_1 \perp P_2$ is called the *topological mating* of the polynomials P_1, P_2 . Moreover, if $\mathcal{K}(P_1) \perp \mathcal{K}(P_2)$ is homeomorphic to a 2-sphere, we say that the topological mating is Moore-unobstructed. We refer the reader to [PM12, Theorem 2.12] for the statement of Moore's theorem, which provides a general sufficient condition for the quotient of \mathbb{S}^2 under an equivalence relation to be a topological 2-sphere, and to [PM12, Proposition 4.12] for a useful application of Moore's theorem giving a sufficient condition for the topological mating of P_1, P_2 (as above) to be Moore-unobstructed (note that the conditions of Moore's theorem are not necessary, see [BM17, Example 13.18]). By [PM12, Proposition 4.3], if the topological mating of P_1, P_2 is not Moore obstructed (i.e., if $\mathcal{K}(P_1) \perp \mathcal{K}(P_2) \cong$ \mathbb{S}^2), then $P_1 \perp P_2$ is topologically conjugate to an orientation-preserving branched covering of \mathbb{S}^2 . The following definition relates the topological mating to rational maps of $\widehat{\mathbb{C}}$. We refer the reader to [DH93] for the notion of *Thurston equivalence* appearing below.

Definition 24 [PM12, Definition 4.4] Let the topological mating of $P_1 \perp P_2$ be Moore-unobstructed, and $h : \mathcal{K}(P_1) \perp \mathcal{K}(P_2) \to \mathbb{S}^2$ be a homeomorphism.

- 1. The polynomials P_1, P_2 are called *combinatorially mateable* if they are postcritically finite and if the branched covering $h \circ P_1 \perp P_2 \circ h^{-1} : \mathbb{S}^2 \to \mathbb{S}^2$ is Thurston equivalent to a rational map *R*.
- 2. The polynomials P_1, P_2 are called *conformally/geometrically mateable* if the homeomorphism *h* can be so chosen that $R = h \circ P_1 \perp P_2 \circ h^{-1} : \mathbb{S}^2 \to \mathbb{S}^2$ is a rational map and *h* is conformal on the interior of $\mathcal{K}(P_1) \perp \mathcal{K}(P_2)$.

Conversely, a rational map *R* is said to be combinatorially (respectively, conformally) a mating if there exist polynomials P_1, P_2 satisfying the corresponding property above with $R = h \circ P_1 \perp P_2 \circ h^{-1}$.

The following equivalent definition of conformal mating is often useful in practice (see [PM12, §4.7] for other definitions). In fact, this definition can be easily adapted for the other frameworks of combination theorems that we will discuss in this section (compare Definition 30).

Definition 25 [PM12, Definition 4.14] A rational map $R : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ of degree $d \ge 2$ is said to be the *conformal mating* of two degree *d* monic, centered, polynomials P_1 and P_2 with connected and locally connected filled Julia sets if and only if there exist continuous maps

$$\psi_1 : \mathcal{K}(P_1) \to \widehat{\mathbb{C}} \text{ and } \psi_2 : \mathcal{K}(P_2) \to \widehat{\mathbb{C}},$$

conformal on $int \mathcal{K}(P_1)$, $int \mathcal{K}(P_2)$, respectively, such that

- 1. $\psi_1(\mathcal{K}(P_1)) \cup \psi_2(\mathcal{K}(P_2)) = \widehat{\mathbb{C}},$
- 2. $\psi_i \circ P_i = R \circ \psi_i$, for $i \in \{1, 2\}$, and
- 3. $\psi_1(z) = \psi_2(w)$ if and only if $z \sim w$, where \sim is the equivalence relation defined above.

With the above notions of mating in place, we can now mention the first major results on mateability of complex polynomials. In fact, these provided the first main application of Thurston's theorem on topological characterization for rational maps. The following theorem completely answers the question of conformal mateability of postcritically finite quadratic polynomials (see [DH85, DH84] for a detailed study of the Mandelbrot set, or [Mil00] for a quick introduction).

Theorem 31 [*Ree92*, *Tan92*, *Shi00*] Let $P_1(z) = z^2 + c_1$ and $P_2(z) = z^2 + c_2$ be two postcritically finite quadratic polynomials. Then P_1 and P_2 are conformally mateable if and only if c_1 and c_2 do not belong to conjugate limbs of the Mandelbrot set.

Among other early works on matings of postcritically finite polynomials, we ought to mention the work of Shishikura and Tan which highlighted additional complexities that are absent in the quadratic setting, but arise for cubic rational maps [SL00].

In [Tan97], Tan described the dynamics of postcritically finite cubic Newton maps (these maps, which are obtained by plugging in complex polynomials in Newton's classical root-finding method, form an important and well-studied class of rational maps), and exhibited in the process the fact that a large subclass of such maps are matings. (See also the more recent work [AR16] for a description of certain postcritically infinite cubic Newton maps as matings.)

The next theorem, due to Yampolsky and Zakeri, was the first existence result for conformal matings of polynomials that are not 'close cousins' of postcritically finite ones. A quadratic polynomial *P* is said to have a *bounded type Siegel fixed point* if it has a fixed point z_0 with $P'(z_0) = e^{2\pi i \theta}$ such that the continued fraction expansion of $\theta \in \mathbb{R}/\mathbb{Z}$ has uniformly bounded partial fractions.

Theorem 32 [*YZ01*] Suppose P_1 , P_2 are quadratic polynomials which are not antiholomorphically conjugate and each of which has a bounded type Siegel fixed point. Then P_1 and P_2 are conformally mateable.

The question of unmating a rational map; i.e., deciding whether a given rational map appears as the mating of two polynomials (and if so, whether such a decomposition is unique) has also been studied by various authors. For a general combinatorial

characterization of hyperbolic, postcritically finite rational maps arising as matings, see [Mey14, Theorem 4.2]. The situation is a bit more subtle for non-hyperbolic rational maps, as discussed in the same paper. However, the next theorem gives a positive answer to the unmating question for a class of postcritically finite rational maps:

Theorem 33 [Mey09, Theorem 1.1] Let $R : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ be a postcritically finite rational map such that its Julia set is the whole sphere. Then every sufficiently high iterate $R^{\circ n}$ of R arises as a mating (i.e., is topologically conjugate to the topological mating of two polynomials).

We refer the reader to the excellent survey article [Mey14] for more on this topic.

Matings of geometrically finite polynomials (i.e., a polynomial whose postcritical set intersects the Julia set in a finite set, or equivalently, if every critical point is either preperiodic, or attracted to an attracting or parabolic cycle) were studied by Haïssinsky and Tan using techniques of David homeomorphisms. They showed that two geometrically finite polynomials P_1 and P_2 with connected Julia sets and parabolic periodic points are mateable if and only if the postcritically finite polynomials $\mathcal{T}(P_1), \mathcal{T}(P_2)$ canonically associated to P_1, P_2 (such that $\mathcal{T}(P_i)$ and P_i have topologically conjugate Julia set dynamics, i = 1, 2) are mateable [HT04, Theorem D] (cf. [LMMN20, Theorem 5.2]).

Mating of anti-holomorphic polynomials. Let us now mention a class of antiholomorphic polynomials (anti-polynomials for short) for which a complete solution to the conformal mating problem is known. These are the so-called *critically fixed* anti-polynomials; i.e., anti-polynomials that fix all of their critical points. The proof of the following theorem crucially uses [PL98, Theorem 3.2], which in many situations, facilitates the application of Thurston's topological characterization of rational maps.

Theorem 34 [LLM20, Theorem 1.3] Let P_1 and P_2 be two (marked) anti-polynomials of equal degree $d \ge 2$, where P_1 is critically fixed and P_2 is postcritically finite, hyperbolic. Then there is an anti-rational map R that is the conformal mating of P_1 and P_2 if and only if there is no Moore obstruction.

In the opposite direction, the question of unmating critically fixed anti-rational maps was also settled in [LLM20, Theorem 1.2], and examples of *shared matings* were demonstrated (cf. [Ree10]).

Remark 1) Combined with [LLM20, Lemma 4.20], Theorem 34 yields an effective procedure to decide conformal mateability of a critically fixed anti-polynomial P_1 and a postcritically finite, hyperbolic anti-polynomial P_2 . This is particularly useful in applying Theorem 45 below to concrete examples.

2) It is worth mentioning that the above mating (respectively, unmating) results for critically fixed anti-polynomials (respectively, anti-rational maps) serve as a precise philosophical counterpart of the double limit theorem for Kleinian reflection groups in the complex dynamics world (see the discussion before [LLM20, Theorem 1.3] and [LLM20, §4.3]).

To conclude, we list a few relevant works that we did not touch upon in this survey: [MT12, ES12, Ché12, CPT12, BEK12a, Sha19]. A good part of the mating theory discussed above carries over to the setting of Thurston maps (i.e., postcritically finite, orientation-preserving branched coverings of \mathbb{S}^2), for which we encourage the reader to consult [BM17, BD18]. Several beautiful visual illustrations of polynomial matings can be found in [Ché]. For a list of open questions on polynomial matings, we refer the reader to [BEK⁺12b].

8 Combining rational maps and Kleinian groups

In this Section, we will expound recently developed frameworks for combining polynomials (respectively, anti-polynomials) with Kleinian (respectively, reflection) groups.

8.1 Mating anti-polynomials with reflection groups

Following [LMM20, LMMN20], we introduce in Subsection 8.1.1 a class of Kleinian reflection groups (called *necklace reflection groups*) central to the mating construction, and associate a map (called the *Nielsen map*) to each necklace reflection group that is *orbit equivalent* to the group. In Subsection 8.1.2, we formalize the notion of conformal mating of a necklace group and an anti-polynomial. It turns out that such conformal matings are realized as *Schwarz reflection maps* associated with *quadrature domains*; these objects are defined in Subsection 8.1.3. Subsection 8.1.4 summarizes some of the main results of [LLMM18a, LLMM18b, LMM20], where various explicit examples of Schwarz reflection maps were shown to be conformal matings of necklace groups and anti-polynomials. Finally in Subsection 8.1.5, we state a general combination theorem for necklace groups and anti-polynomials proved in [LMMN20].

8.1.1 Necklace reflection groups

A circle packing is a connected collection of oriented circles in \mathbb{C} with disjoint interiors (where the interior is determined by the orientation). Up to a Möbius map, we can always assume that no circle of the circle packing contains ∞ in its interior; i.e., the interior of each circle *C* of the circle packing can be assumed to be the bounded complementary component int*C* of *C*. Combinatorially, a circle packing can be described by its *contact graph*, where we associate a vertex to each circle, and connect two vertices by an edge if and only if the two associated circles touch. By the Koebe-Andreev-Thurston circle packing theorem [Thu80, Corollary 13.6.2], every connected, simple, planar graph is the contact graph of some circle packing.

Definition 26 A *necklace reflection group* is a group generated by reflections in the circles of a finite circle packing whose contact graph is 2-*connected* and *outerplanar*; i.e., the contact graph remains connected if any vertex is deleted, and has a face containing all the vertices on its boundary.

Note that since a necklace reflection group is a discrete subgroup of the group of all Möbius and anti-Möbius automorphisms of $\widehat{\mathbb{C}}$, definitions of limit set and domain of discontinuity can be easily extended to necklace reflection groups. By [LLM20, Proposition 3.4], the limit set of a necklace reflection group is connected. Moreover, for a necklace reflection group Γ generated by reflections in the circles C_1, \dots, C_{d+1} , the set

$$\mathcal{F}_{\Gamma} := \widehat{\mathbb{C}} \setminus \left(\bigcup_{i=1}^{d+1} \operatorname{int} C_i \bigcup_{j \neq k} (C_j \cap C_k) \right)$$

is a fundamental domain for the Γ -action on $\Omega(\Gamma)$ [LMM20, Proposition 2.14].

To a necklace reflection group Γ , one can associate a piecewise anti-Möbius reflection map ρ_{Γ} that plays an important role in the mating construction.

Definition 27 [LMM20, Definition 2.29], [LMMN20, Definition 6.6] Let Γ be a necklace reflection group generated by reflections $\{r_i\}_{i=1}^{d+1}$ in circles $\{C_i\}_{i=1}^{d+1}$. We define the associated *Nielsen map* ρ_{Γ} by:

$$\rho_{\Gamma}: \bigcup_{i=1}^{d+1} \overline{\operatorname{int} C_i} \to \widehat{\mathbb{C}}, \quad z \longmapsto r_i(z) \text{ if } z \in \overline{\operatorname{int} C_i}.$$

The next proposition underscores the intimate dynamical connection between a necklace group Γ and its Nielsen map ρ_{Γ} .

Proposition [LMM20, Proposition 2.31] Let Γ be a necklace reflection group. The map ρ_{Γ} is orbit equivalent to Γ on $\widehat{\mathbb{C}}$; i.e., for any two points $z, w \in \widehat{\mathbb{C}}$, there exists $g \in \Gamma$ with g(z) = w if and only if there exist non-negative integers n_1, n_2 such that $\rho_{\Gamma}^{\circ n_1}(z) = \rho_{\Gamma}^{\circ n_2}(w)$.

The simplest examples of necklace reflection groups are regular ideal polygon reflection groups.

Definition 28 Consider the Euclidean circles C_1, \dots, C_{d+1} where C_j intersects \mathbb{S}^1 at right angles at the roots of unity $\exp(\frac{2\pi i \cdot (j-1)}{d+1})$, $\exp(\frac{2\pi i \cdot j}{d+1})$. (By [VS93, Part II, Chapter 5, Theorem 1.2], the group generated by reflections in these circles is discrete.) We denote this group by Γ_{d+1} .

Note that $\rho_{\Gamma_{d+1}}$ restricts to an expansive degree *d* orientation-reversing covering of \mathbb{S}^1 . By [CR80], there exists a homeomorphism \mathcal{E}_d of the circle that conjugates $\rho_{\Gamma_{d+1}}$ to \overline{z}^d . The conjugacy \mathcal{E}_d serves as a connecting link between reflection groups and quadratic anti-polynomials.

8.1.2 Conformal mating of anti-polynomials and necklace groups

The precise meaning of conformal matings of the Nielsen map of a necklace group and an anti-polynomial is given below. The definition is an adaptation of the classical definition of conformal matings of two polynomials.

Let Γ be a necklace group generated by reflections in circles C_1, \dots, C_{d+1} . The unbounded component of the domain of discontinuity $\Omega(\Gamma)$ is Γ -invariant [LMM20, Proposition 2.28], and we denote it by $\Omega_{\infty}(\Gamma)$. We also set $\mathcal{K}(\Gamma) := \mathbb{C} \setminus \Omega_{\infty}(\Gamma)$. According to [LMM20, Proposition 2.36], the restriction of ρ_{Γ} to $\Omega_{\infty}(\Gamma)$ is conformally conjugate to the $\rho_{\Gamma_{d+1}}$ -action on $\mathbb{C} \setminus \overline{\mathbb{D}}$, and (the inverse of) this conformal conjugacy continuously extends to yield a semiconjugacy $\phi_{\Gamma} : \mathbb{S}^1 \to \Lambda(\Gamma) = \partial \mathcal{K}(\Gamma)$ between $\rho_{\Gamma_{d+1}}|_{\mathbb{S}^1}$ and $\rho_{\Gamma}|_{\Lambda(\Gamma)}$ such that $\phi_{\Gamma}(1)$ is the point of tangential intersection of C_1 and C_{d+1} . Recall also that $\mathcal{E}_d : \mathbb{S}^1 \to \mathbb{S}^1$ is a topological conjugacy between $\rho_{\Gamma_{d+1}}|_{\mathbb{S}^1}$.

Let *P* be a monic, centered, anti-polynomial of degree *d* such that $\mathcal{J}(P)$ is connected and locally connected. Denote by $\phi_P : \mathbb{D}^* \to \mathcal{B}_{\infty}(P)$ the Böttcher coordinate for *P* such that $\phi'_P(\infty) = 1$. We note that since $\partial \mathcal{K}(P) = \mathcal{J}(P)$ is locally connected by assumption, it follows that ϕ_P extends to a semiconjugacy between $z \mapsto \overline{z}^d|_{\mathbb{S}^1}$ and $P|_{\mathcal{J}(P)}$.

The equivalence relation below specifies a gluing of $\mathcal{K}(\Gamma)$ with $\mathcal{K}(P)$ along their boundaries. The presence of the topological conjugacy \mathcal{E}_d in the definition of the equivalence relation ensures that the maps ρ_{Γ} and P fit together to produce a continuous map on the resulting topological 2-sphere (when there is no Moore obstruction).

Definition 29 We define the equivalence relation ~ on $\mathcal{K}(\Gamma) \sqcup \mathcal{K}(P)$ generated by $\phi_{\Gamma}(t) \sim \phi_{P}(\overline{\mathcal{E}_{d}(t)})$ for all $t \in \mathbb{S}^{1}$.

The following definition essentially says that an anti-holomorphic map F (defined on a subset of the Riemann sphere) is a conformal mating of Γ and P if there are continuous semi-conjugacies from $\mathcal{K}(\Gamma), \mathcal{K}(P)$ into the dynamical plane of F(conformal on the interiors) such that the images fill up the whole sphere and intersect only along their boundaries as prescribed by the equivalence relation ~ (compare Definition 25).

Definition 30 [LMMN20, Definition 10.16] Let Γ be a necklace group as above, and let *P* be a monic, centered anti-polynomial such that $\mathcal{J}(P)$ is connected and locally connected. Further, let $\Omega \subsetneq \widehat{\mathbb{C}}$ be an open set, and $F : \overline{\Omega} \to \widehat{\mathbb{C}}$ be a continuous map that is anti-meromorphic on Ω . We say that *F* is a *conformal mating* of Γ with *P* if there exist continuous maps

$$\psi_P : \mathcal{K}(P) \to \widehat{\mathbb{C}} \text{ and } \psi_\Gamma : \mathcal{K}(\Gamma) \to \widehat{\mathbb{C}},$$

conformal on $int\mathcal{K}(P)$, $int\mathcal{K}(\Gamma)$, respectively, such that

1. $\psi_P(\mathcal{K}(P)) \bigcup \psi_{\Gamma}(\mathcal{K}(\Gamma)) = \widehat{\mathbb{C}},$

2. $\Omega = \widehat{\mathbb{C}} \setminus \psi_{\Gamma}(\overline{\mathcal{F}_{\Gamma}}),$ 3. $\psi_{P} \circ P = F \circ \psi_{P} \text{ on } \mathcal{K}(P),$ 4. $\psi_{\Gamma} \circ \rho_{\Gamma} = F \circ \psi_{\Gamma} \text{ on } \mathcal{K}(\Gamma) \setminus \text{int}\mathcal{F}_{\Gamma}, \text{ and}$ 5. $\psi_{\Gamma}(z) = \psi_{P}(w) \text{ if and only if } z \sim w \text{ where } \sim \text{ is as in Definition 29}.$

Remark For the purposes of mating necklace groups with anti-polynomials, it is important to work with labeled circle packings, or equivalently, to regard the space of necklace groups as a space of representations of the ideal polygon reflection group Γ_{d+1} . While we have suppressed this abstraction for ease of exposition, we refer the reader to [LMM20, §2.2] or [LMMN20, §10.1], where necklace groups are organized in Bers slices of Γ_{d+1} . Although this point of view may seem like an artificial complication at a first glance, the language of representations turns out to be an unavoidable technicality in the mating theory. Roughly speaking, different representations give rise to different ways of gluing the limit set of a necklace group with the Julia set of an anti-polynomial, and the choice of gluing determines whether or not a conformal mating exists (compare [LMMN20, Remark 10.21]).

8.1.3 Schwarz reflection maps

By definition, a domain $\Omega \subseteq \widehat{\mathbb{C}}$ satisfying $\infty \notin \partial \Omega$ and $\Omega = \operatorname{int}\overline{\Omega}$ is a *quadrature domain* if there exists a continuous function $\sigma : \overline{\Omega} \to \widehat{\mathbb{C}}$ such that σ is antimeromorphic in Ω and $\sigma(z) = z$ on the boundary $\partial \Omega$. Such a function σ is unique (if it exists), and is called the *Schwarz reflection map* associated with Ω . (See [AS76], [LM16] and the references therein.)

It is well known that except for a finite number of *singular* points (cusps and double points), the boundary of a quadrature domain consists of finitely many disjoint real analytic curves [Sak91]. Every non-singular boundary point has a neighborhood where the local reflection in $\partial\Omega$ is well-defined. The (global) Schwarz reflection σ is an anti-holomorphic continuation of all such local reflections.

Round disks on the Riemann sphere are the simplest examples of quadrature domains. Their Schwarz reflections are just the usual circle reflections. Further examples can be constructed using univalent polynomials or rational functions. In fact, simply connected quadrature domains admit a simple characterization.

Proposition [AS76, Theorem 1] A simply connected domain $\Omega \subseteq \widehat{\mathbb{C}}$ with $\infty \notin \partial \Omega$ and $\operatorname{int} \overline{\Omega} = \Omega$ is a quadrature domain if and only if the Riemann uniformization $f : \mathbb{D} \to \Omega$ extends to a rational map on $\widehat{\mathbb{C}}$.

In this case, the Schwarz reflection map σ of Ω is given by $f \circ (1/\overline{z}) \circ (f|_{\mathbb{D}})^{-1}$. Moreover, if the degree of the rational map f is d, then $\sigma : \sigma^{-1}(\Omega) \to \Omega$ is a (branched) covering of degree (d-1), and $\sigma : \sigma^{-1}(\operatorname{int}\Omega^c) \to \operatorname{int}\Omega^c$ is a (branched) covering of degree d.

$$\begin{array}{c} \overline{\mathbb{D}} & \stackrel{\phi}{\longrightarrow} \overline{\Omega} \\ 1/\overline{z} \downarrow & \qquad \downarrow^{c} \\ \widehat{\mathbb{C}} \setminus \mathbb{D} & \stackrel{\phi}{\longrightarrow} \widehat{\mathbb{C}}. \end{array}$$

In [LM16], questions on equilibrium states of certain 2-dimensional Coulomb gas models were answered using iteration of Schwarz reflection maps associated with quadrature domains. It transpired from their work that these maps give rise to dynamical systems that are interesting in their own right. The general situation is as follows. Given a disjoint collection of quadrature domains, we call the complement of their union a *droplet*. Removing the double points and cusps from the boundary of a droplet yields the *desingularized droplet* or the *fundamental tile*. One can then look at a partially defined anti-holomorphic dynamical system σ that acts on (the closure of) each quadrature domain as its Schwarz reflection map. Under this dynamical system, the Riemann sphere \mathbb{C} admits a dynamically invariant partition. The first one is an open set called the escaping/tiling set, it is the set of all points that eventually escape to the fundamental tile (on the interior of which σ is not defined). The second invariant set is the *non-escaping* set, the complement of the tiling set or equivalently, the set of all points on which σ can be iterated forever. When the tiling set contains no critical points of σ , it is often the case that the dynamics of σ on its non-escaping set resembles that of an anti-polynomial on its filled Julia set, while the σ -action on the tiling set exhibits features of reflection groups.

8.1.4 Examples of the mating phenomenon

By studying the dynamics and parameter spaces of specific families of Schwarz reflection maps, one can often recognize such maps as matings of anti-polynomials and necklace reflection groups. This strategy was successfully implemented in [LLMM18a, LLMM18b, LMM20]. We collect some results from these papers in this subsection.

Example 1: The deltoid reflection. We will start with the simplest instance of the mating phenomenon; namely, the conformal mating of the anti-polynomial \overline{z}^2 and the ideal triangle reflection group Γ_3 .

Theorem 39 [*LLMM18a*, Theorem 1.1] The map $f_0(z) = 1/z + z^2/2$ is injective on $\overline{\mathbb{D}}$, and hence $\Omega_0 := f_0(\mathbb{D})$ is a simply connected quadrature domain. The associated Schwarz reflection map σ_0 is the unique conformal mating of \overline{z}^2 and Γ_3 .

Remark A welding homeomorphism is a homeomorphism of the circle that arises as the composition of a conformal map from the unit disk onto the interior region of a Jordan curve with a conformal map from the exterior of this Jordan curve onto the exterior of the unit disk. A complex-analytic corollary of Theorem 39 is that the circle homeomorphisms \mathcal{E}_2 is a welding homeomorphism. That the same is true for each \mathcal{E}_d ($d \ge 2$) follows from a straightforward higher degree generalization



Fig. 2 Left: The tessellation of \mathbb{D} for the ideal triangle reflection group. Right: The dynamical plane of the Schwarz reflection map associated with $\Omega_0 = f_0(\mathbb{D})$ (the exterior of a deltoid curve). The dynamics on the exterior of the bright green fractal curve is conformally conjugate to $\overline{z}^2|_{\mathbb{D}}$, while the dynamics on the interior is conformally equivalent to the Nielsen map of Γ_3 .

of Theorem 39 worked out in [LLMM19, Appendix B] (also compare Theorem 44 below). We refer the reader to [LMMN20, Theorem 5.1] for a general conformal welding result for circle homeomorphisms conjugating suitable covering maps of the circle.

Example 2: The circle and cardioid family. To describe Schwarz reflection maps that are conformal matings of other quadratic anti-polynomials with the ideal triangle reflection group, we need to recall the Circle and Cardioid family which was introduced in [LLMM18a, §6]. We consider the fixed cardioid

$$\heartsuit := \{ w = z/2 - z^2/4 : |z| < 1 \},\$$

and for each complex number $a \in \mathbb{C} \setminus (-\infty, -1/12)$, let $B(a, r_a)$ be the smallest open disk containing \heartsuit centered at *a* (in other words, $\{w : |w - a| = r_a\}$ is a circumcircle of the cardioid; see [LLMM18b, Figure 2]). Let $\Omega_a := \heartsuit \cup \overline{B}(a, r_a)^c$ (where $\overline{B}(a, r_a)$ is the closed disk $\{w : |w - a| \le r_a\}$), and $T_a := \Omega_a^c$. We now define a piecewise Schwarz reflection dynamical system $F_a : \overline{\Omega}_a \to \widehat{\mathbb{C}}$ as,

$$w \mapsto \begin{cases} \sigma(w) & \text{if } w \in \overline{\heartsuit}, \\ \sigma_a(w) & \text{if } w \in B(a, r_a)^c, \end{cases}$$

where σ is the Schwarz reflection of \heartsuit , and σ_a is reflection with respect to the circle $\partial B(a, r_a)$. The family

$$\mathcal{S} := \left\{ F_a : \overline{\Omega}_a \to \widehat{\mathbb{C}} : a \in \mathbb{C} \setminus (-\infty, -1/12) \right\}$$

is referred to as the C&C family.

For any $a \in \mathbb{C} \setminus (-\infty, -1/12)$, ∂T_a has two singular points; namely, the double point α_a where $\partial B(a, r_a)$ touches $\partial \heartsuit$, and the cusp point $\frac{1}{4}$. Both of them are fixed points of F_a . The *fundamental tile* of F_a is defined as $T_a^0 := T_a \setminus \{\alpha_a, \frac{1}{4}\}$. A parameter $a \in \mathbb{C} \setminus (-\infty, -1/12)$ (equivalently, the corresponding map $F_a \in S$) is said to be *postcritically finite* if the unique (simple) critical point 0 of F_a has a finite forward orbit that does not meet T_a^0 . The following mating description for postcritically finite maps in S was given in [LLMM18b].



Fig. 3 Under the bijection χ of Theorem 41, the postcritically finite quadratic anti-polynomial $\overline{z}^2 - 1$ corresponds to F_a with a = 0. Left: The filled Julia set of $\overline{z}^2 - 1$. Right: The part of the non-escaping set of F_0 inside the cardioid (in dark blue) with the critical point 0 marked. Both maps have a critical cycle of period 2.

Theorem 41 [LLMM18a, §8], [LLMM18b, Theorems 1.1, 1.2] There exists a bijection χ between postcritically finite maps in S and (the Möbius conjugacy classes of) postcritically finite quadratic anti-polynomials $\overline{z}^2 + c$ (excluding \overline{z}^2) such that the postcritically finite map $F_a \in S$ is a conformal mating of the ideal triangle reflection group Γ_3 and the quadratic anti-polynomial $\overline{z}^2 + \chi(a)$.

Remark For conformal matings of Γ_3 with more general quadratic anti-polynomials, and combinatorial relations between the *connectedness loci* of quadratic anti-polynomials and the family S, see [LLMM18b].

Example 3: The space Σ_d^* . The family of 'univalent rational maps'

$$\Sigma_d^* := \left\{ g(z) = z + \frac{a_1}{z} + \dots + \frac{a_d}{z^d} : a_d = -\frac{1}{d} \text{ and } g|_{\widehat{\mathbb{C}}\setminus\overline{\mathbb{D}}} \text{ is conformal} \right\}$$

was introduced in [LM14] and studied extensively in [LMM19] in terms of the associated Schwarz reflection maps.

Remark The family Σ_d^* is closely related to the classically studied space Σ of suitably normalized schlicht functions on $\widehat{\mathbb{C}} \setminus \overline{\mathbb{D}}$, see [Dur83, §4.7, §9.6].

Combining the pinching deformation theory for Σ_d^* (developed in [LMM19]) with tools from holomorphic dynamics, it was proved in [LMM20] that:



Fig. 4 Left: The dynamical plane of the Schwarz reflection map associated with some $f \in \partial \Sigma_4^*$. Right: The limit set of the corresponding necklace reflection group Γ_f .

Theorem 44 [LMM20, Theorem A] There is a bijection $f \mapsto \Gamma_f$ between Σ_d^* and the space of necklace reflection groups of rank d + 1 (up to a natural equivalence) such that the Schwarz reflection map associated with $f \in \Sigma_d^*$ is a conformal mating of the anti-polynomial \overline{z}^d with the corresponding necklace group Γ_f .

8.1.5 The general theorem

We conclude our discussion of combinations of necklace reflection groups and antipolynomials with a general existence theorem:

Theorem 45 [LMMN20, Lemma 10.17, Theorem 10.20] Let P be a monic, postcritically finite, hyperbolic anti-polynomial of degree d, and let Γ be a necklace group. Then, P and Γ are conformally mateable if and only if $\mathcal{K}(P) \sqcup \mathcal{K}(\Gamma)/\sim$ is homeomorphic to \mathbb{S}^2 (where \sim is the equivalence relation from Definition 29).

Moreover, if $F : \overline{\Omega} \to \widehat{\mathbb{C}}$ is a conformal mating of Γ and P, then each component of Ω is a simply connected quadrature domain, and F is the piecewise defined Schwarz reflection map associated with these quadrature domains.

The hard part of the above theorem is to show that if $\mathcal{K}(P) \sqcup \mathcal{K}(\Gamma)/\sim$ is homeomorphic to \mathbb{S}^2 , then a conformal mating of *P* and Γ exists. In fact, the condition that $\mathcal{K}(P) \sqcup \mathcal{K}(\Gamma)/\sim \cong \mathbb{S}^2$ guarantees the existence of a topological mating on a 2-sphere, but promoting the topological mating to an anti-holomorphic map lies at the heart of the difficulty. This goal is achieved in two steps. One first uses Thurston's topological characterization theorem to construct a hyperbolic anti-rational map *R* that is a conformal mating of *P* and another postcritically finite (in fact, critically fixed),





Fig. 5 Bottom left: The circles C_i generate a necklace reflection group Γ . Bottom right: The dynamical plane of $P(z) = \overline{z}^3 - \frac{3i}{\sqrt{2}}\overline{z}$; each critical point of which forms a 2-cycle. Top: The conformal mating of P and the necklace group Γ is given by the piecewise Schwarz reflection map associated with the disjoint union of three quadrature domains: the exterior of an ellipse, and two round disks contained in the interior of the ellipse. Each of the two critical points of F forms a 2-cycle. (See [LMMN20, §11.2] for proofs of these statements.)

hyperbolic anti-polynomial P_{Γ} such that the Julia dynamics of P_{Γ} is topologically conjugate to the limit set dynamics of the Nielsen map ρ_{Γ} . The existence of such an anti-polynomial P_{Γ} follows from [LMM20] or [LLM20], while conformal mateability of *P* and P_{Γ} follows from the general mateability criterion given in Theorem 34 (in fact, the condition $\mathcal{K}(P) \sqcup \mathcal{K}(\Gamma)/\sim \cong S^2$ is equivalent to saying that the topological mating of *P* and P_{Γ} is Moore-unobstructed, so Theorem 34 can be applied to produce *R*). Finally, to turn *R* into a conformal mating of *P* and Γ , one needs to glue Nielsen maps of ideal polygon reflection groups in suitable invariant Fatou components of *R*. The fact that all fixed points of *R* on its Julia set are hyperbolic while those of a

Nielsen map are parabolic prohibits the use of purely quasiconformal tools to carry out this task. This problem is tackled by employing surgery techniques involving David homeomorphisms: generalizations of quasiconformal homeomorphisms.

Remark Although Theorem 45 guarantees the existence of conformal matings of suitable anti-polynomials and necklace reflection groups, in general, it may be hard to find explicit Schwarz reflection maps realizing such conformal matings. However, in certain low complexity situations, the second statement of the theorem (that the conformal matings are piecewise Schwarz reflection maps associated with simply connected quadrature domains) allows one to use Proposition 38 and the desired dynamical properties to explicitly characterize the conformal matings (see Figure 5 for an illustration, and [LMMN20, §11] for various worked out examples).

8.2 Mating polynomials with Kleinian groups

This subsection is a summary of [MM21], where a new setup for combination theorems of complex polynomials and Kleinian surface groups was designed using the notion of orbit equivalence.

8.2.1 The Fuchsian case

A foundational problem that arises in trying to make sense of what it means to combine a polynomial P with a Kleinian group Γ is that on one side of the picture we have the semigroup $\langle P \rangle$ generated by P, while on the other side we have a noncommutative group Γ generated by more than one element. To formulate a precise notion of mateability between Fuchsian groups and complex polynomials (with Jordan curve Julia sets), one needs to address this inherent discord between these two objects, and this leads to the notion of *mateable* circle maps: *single* maps A that capture essential dynamical and combinatorial features of Fuchsian groups acting on \mathbb{S}^1 . Further, A should also be dynamically compatible with polynomial maps. Before giving a precise definition of mateable maps, let us outline the underlying motivation: the following features are required of a mateable map $A : \mathbb{S}^1 \to \mathbb{S}^1$.

- 1. A must be dynamically compatible with a Fuchsian group Γ . This leads to
 - a. orbit equivalence between A and Γ .
 - b. A has to be piecewise Fuchsian.
- 2. *A* must be dynamically compatible with complex polynomials. Hence we demand the existence of a topological conjugacy between *A* and the polynomial $z^d|_{\mathbb{S}^1}$ (where $d \ge 2$ is the degree of *A*),
- 3. A must be combinatorially compatible with z^d leading to a Markov condition,

4. A must be conformally compatible with z^d requiring absence of asymmetrically hyperbolic periodic break-points of A (this is a weaker version of the C^1 -condition).

To fulfill the above requirements, we make the following definition.

Definition 31 [MM21, Definitions 2.7, 2.16]

1. A map $A : \mathbb{S}^1 \to \mathbb{S}^1$ is called *piecewise Möbius* if there exist $k \in \mathbb{N}$, closed arcs $I_j \subset \mathbb{S}^1$, and $g_j \in \operatorname{Aut}(\mathbb{D}), j \in \{1, \dots, k\}$, such that

a.
$$\mathbb{S}^1 = \bigcup_{j=1}^k I_j$$
,
b. $\operatorname{int} I_m \cap \operatorname{int} I_n = \emptyset$ for $m \neq n$, and
c. $A|_{I_j} = g_j$.

A piecewise Möbius map is called *piecewise Fuchsian* if g_1, \dots, g_k generate a Fuchsian group, which we denote by Γ_A .

- 2. A map $A : \mathbb{S}^1 \to \mathbb{S}^1$ is called *piecewise Fuchsian Markov* if it is a piecewise Fuchsian expansive covering map (of degree at least two) such that the pieces (intervals of definition) of A form a Markov partition for $A : \mathbb{S}^1 \to \mathbb{S}^1$.
- 3. A piecewise Fuchsian Markov map *A* is said to be *mateable* if *A* is orbit equivalent to the Fuchsian group Γ_A generated by its pieces, and none of the periodic breakpoints of *A* is asymmetrically hyperbolic.

We refer the reader to [MM21, §2] for the definition of the term 'symmetrically hyperbolic', and for a detailed discussion on the necessity of each of the requirements in the definition of a mateable map. We also note that the expansivity condition above ensures that a mateable map is topologically conjugate to the polynomial z^d (for some $d \ge 2$).

Remark In the anti-holomorphic setting, the role of mateable maps was played by Nielsen maps of necklace reflection groups (see Subsection 8.1.1). \Box

The simplest example of a mateable map is given by the classical *Bowen-Series map* [Bow79, BS79]. While such a map can be defined for arbitrary Fuchsian groups equipped with suitable fundamental domains, they are typically discontinuous. However, it turns out that for Fuchsian groups uniformizing spheres with punctures (possibly with one/two order two orbifold points), the Bowen-Series map is a covering map of the circle satisfying the defining properties of a mateable map [MM21, §2].

Higher Bowen-Series maps. More examples of mateable maps are given by *higher Bowen-Series maps* of punctured sphere Fuchsian groups (see [MM21, §4] for their definition and basic properties). As suggested by the name, there are close connections between higher Bowen-Series maps and Bowen-Series maps. Indeed, the higher Bowen-Series map of a Fuchsian group uniformizing $S_{0,k}$ (a sphere with *k* punctures) can be represented as the second iterate of the Bowen-Series map of a Fuchsian group uniformizing a sphere with roughly k/2 punctures and zero/one/two



Fig. 6 Left: *R* is a fundamental domain of a Fuchsian group Γ uniformizing $S_{0,3}$ with side pairing transformations $g_1^{\pm 1}, g_2^{\pm 1}$. The canonical extension of the Bowen-Series map of Γ (associated with *R*) is defined on $\overline{\mathbb{D}} \setminus \operatorname{int} R$ in terms of $g_1^{\pm 1}, g_2^{\pm 1}$ as shown in the figure. Right: The second iterate of the Bowen-Series map of Γ is a higher Bowen-Series map of the index two subgroup $\Gamma' = \langle g_1^2, g_1g_2, g_1g_2^{-1} \rangle \leq \Gamma$, which uniformizes $S_{0,4}$. Its canonical extension is defined on the region enclosed by \mathbb{S}^1 and the red (hyperbolic) geodesics in terms of $h_1 = g_1^2, h_2 = g_1g_2, h_3 = g_1g_2^{-1}$ as shown in the figure. It maps the boundary of the red polygon onto the boundary of *R*. The degree of the Bowen-Series map of Γ (as a circle covering) is 3, so the degree of the higher Bowen-Series map of Γ' is 9.

order two orbifold points [MM21, Corollary 5.6] (see Figure 6). Alternatively, a higher Bowen-Series map of a Fuchsian group is obtained by 'gluing together' several Bowen-Series maps of the same Fuchsian group with overlapping fundamental domains [MM21, Proposition 4.5] (see Figure 7).

Every piecewise Fuchsian Markov map A of the circle can be conformally extended to a canonically defined subset of $\overline{\mathbb{D}}$ (see [MM21, §2.2]). This extension is termed the *canonical extension of A*. The following result, which is a conformal combination theorem for punctured sphere Fuchsian groups and hyperbolic polynomials with Jordan curve Julia sets, can be regarded as an analog of the Bers' simultaneous uniformization theorem in the current setting.

Theorem 48 [MM21, Theorem 3.7, Theorem 4.8] The canonical extensions of Bowen-Series maps and higher Bowen-Series maps of Fuchsian groups uniformizing punctured spheres (possibly with one/two orbifold points of order two) can be conformally mated with polynomials lying in principal hyperbolic components (of appropriate degree).

Remark As in the anti-holomorphic case, there is a key qualitative difference between the dynamics of Bowen-Series (respectively, higher Bowen-Series) maps on \mathbb{S}^1 and the dynamics of polynomials (lying in principal hyperbolic components) on their Julia set; namely, the former has parabolic fixed points on \mathbb{S}^1 while all fixed points of the latter on their Julia sets are repelling. Consequently, the topological conjugacy between a Bowen-Series (respectively, higher Bowen-Series) map



Fig. 7 The quadrilaterals *D* and *P* with ideal vertices at 1, 2, 3, 4 and 1, 2, 3, 4 (respectively) together form a fundamental domain of a Fuchsian group Γ uniformizing $S_{0,4}$ with side pairing transformations $g_1^{\pm 1}, g_2^{\pm 1}, g_3^{\pm 1}$. The corresponding higher Bowen-Series map *A* of Γ acts on the anti-clockwise arc from 1 to 4 as the Bowen-Series map of Γ associated with the fundamental domain $D \cup P$; while on the clockwise arc from *j* to j + 1, *A* equals the Bowen-Series map of Γ associated with the fundamental domain $D \cup g_j(P)$ (j = 1, 2, 3). The degree of a Bowen-Series map of Γ (associated with any fundamental domain) is 5, while the degree of a higher Bowen-Series map of Γ is 9.

and such a polynomial is not quasisymmetric. This forces one to abandon classical quasiconformal techniques (used in the proof of Bers' simultaneous uniformization theorem), and apply David homeomorphisms to prove Theorem 48. $\hfill \Box$

Moduli space of Fuchsian matings. In the torsion-free case, the only topological surfaces that Theorem 48 succeeds to combine with complex polynomials are punctured spheres (see [MM21, 6.35] for the definition of moduli space of matings between a topological surface and hyperbolic complex polynomials with Jordan curve Julia sets). This naturally raises the following questions.

- 1. Do mateable maps exist for higher genus surfaces (possibly with punctures)? More precisely, does there exist a mateable map A with $\mathbb{D}/\Gamma_A \cong S_{g,k}$, for $g \ge 1$?
- 2. Are Bowen-Series and higher Bowen-Series maps the only mateable maps associated with punctured spheres?

In this generality, the above questions remain open. However, [MM21, Theorems 6.18, 6.33] give a complete description of mateable maps satisfying some natural 2-point conditions over and above orbit equivalence. It turns out that under these additional hypotheses, punctured spheres are the only topological surfaces that can be combined with complex polynomials (see [MM21, Theorem 6.36] for a complete description of the interiors of such constrained moduli space of matings). A major part of the proofs of these theorems is to determine the topology of the surface \mathbb{D}/Γ_A from the dynamical properties of a mateable map *A*, and this is accomplished by analyzing certain patterns and laminations associated with mateable maps.

8.2.2 The case of Bers' boundary groups

We proceed to discuss the structure of the boundaries of the moduli spaces of Fuchsian matings arising from Theorem 48.

For definiteness, let us fix a base Fuchsian group Γ_0 uniformizing $S_{0,k}$ ($k \ge 3$). For each Γ lying on the boundary of the Bers slice $\mathcal{B}(\Gamma_0)$, there exists a continuous map $\phi_{\Gamma} : \mathbb{S}^1 \to \Lambda_{\Gamma}$, called the *Cannon-Thurston map* after [CT07], that semi-conjugates the action of Γ_0 to that of Γ [Mj14a, Mj14b, DM16, Mj17]. In fact, the data of the *ending lamination* can be recovered from the Cannon-Thurston map. More precisely, the group Γ can be obtained by 'pinching a lamination' on the surface \mathbb{D}/Γ_0 (while keeping the hyperbolic structure on the surface $(\widehat{\mathbb{C}} \setminus \overline{\mathbb{D}})/\Gamma_0$ unchanged), and the endpoints of the corresponding geodesic lamination on \mathbb{D} generate a Γ_0 -invariant equivalence relation on \mathbb{S}^1 which agrees with the one defined by the fibers of the Cannon-Thurston map ϕ_{Γ} .

Assume further that A_{Γ_0} is a Bowen-Series (respectively, higher Bowen-Series) map of Γ_0 . To extend the notion of mateability to a group $\Gamma \in \partial \mathcal{B}(\Gamma_0)$, one needs the limit set of Γ to carry a continuous, piecewise complex-analytic self-map A_{Γ} (that is orbit equivalent to Γ) defined by the following commutative diagram:

$$\begin{array}{c} \mathbb{S}^1 \xrightarrow{A_{\Gamma_0}} \mathbb{S}^1 \\ \downarrow^{\phi_{\Gamma}} \downarrow & \downarrow^{\phi_{\Gamma}} \\ \Lambda_{\Gamma} \xrightarrow{A_{\Gamma}} \Lambda_{\Gamma} \end{array}$$

See [MM21, §7.1] for details and an alternative description of A_{Γ} as a uniform limit of Bowen-Series (respectively, higher Bowen-Series) maps. The map A_{Γ} , if it exists, is called the *Bowen-Series* (respectively, *higher Bowen-Series*) map of Γ and can be thought of as a mateable map associated with a Bers' boundary group.

It turns out that the existence of such a map A_{Γ} imposes severe restrictions on the laminations that can be pinched. The next theorem says that only finitely many possibilities exist. We call such laminations *admissible* (see Figure 8 for an example of an admissible lamination in the Bowen-Series case).

For $\Gamma \in \partial \mathcal{B}(\Gamma_0)$, we denote the unique Γ -invariant component of the domain of discontinuity $\Omega(\Gamma)$ by $\Omega_{\infty}(\Gamma)$, and set $K(\Gamma) := \widehat{\mathbb{C}} \setminus \Omega_{\infty}(\Gamma)$. If Γ admits a Bowen-Series (respectively, higher Bowen-Series) map $A_{\Gamma} : \Lambda_{\Gamma} \to \Lambda_{\Gamma}$, then this map can be extended as a continuous, piecewise Möbius map to a canonical closed set $K(\Gamma) \setminus \operatorname{int} R_{\Gamma}$, where R_{Γ} is a 'pinched' fundamental domain for the Γ -action on $\Omega(\Gamma) \setminus \Omega_{\infty}(\Gamma)$ determined by R_{Γ_0} (see [MM21, §7.3]). This canonical extension is denoted by \widehat{A}_{Γ} . The following theorem also demonstrates conformal mateability of groups $\Gamma \in \partial \mathcal{B}(\Gamma_0)$ admitting Bowen-Series/ higher Bowen-Series maps with polynomials lying in principal hyperbolic components (of suitable degree).

Theorem 50 [MM21, Lemma 7.3, Lemma 7.5, Theorem 7.19] Let Γ_0 be a Fuchsian group uniformizing $S_{0,k}$. Then, there are only finitely many quasiconformal conjugacy classes of groups $\Gamma \in \partial \mathcal{B}(\Gamma_0)$ for which the Cannon-Thurston map of Γ semi-



Fig. 8 Left: R_{Γ_0} is a fundamental domain of a Fuchsian group Γ_0 uniformizing $S_{0,7}$ with side pairing transformations $g_1^{\pm 1}, \dots, g_6^{\pm 1}$. The geodesic lamination \mathcal{L}^* on \mathbb{D}/Γ_0 consisting of two simple, closed curves corresponding to the elements $g_5, g_2g_5^{-1} \in \Gamma_0$ is admissible for the Bowen-Series map A_{Γ_0} . The blue and green geodesics are the connected components of the Γ_0 -lift of \mathcal{L}^* that intersect R_{Γ_0} . Right: A cartoon of the limit set of a Bers' boundary group Γ , which is obtained by pinching \mathcal{L}^* . The Γ -action on $\Omega(\Gamma) \setminus \Omega_{\infty}(\Gamma)$ admits a pinched fundamental domain R_{Γ} , and the canonical extension \widehat{A}_{Γ} of the Bowen-Series map of Γ is defined on $K(\Gamma) \setminus \operatorname{int} R_{\Gamma}$. Two of the components of $\Omega(\Gamma)$ intersecting R_{Γ} are invariant under \widehat{A}_{Γ} , while the other two components U^{\pm} form a 2-cycle. The first return map of \widehat{A}_{Γ} on U^{\pm} are conformally conjugate to higher Bowen-Series maps of punctured sphere Fuchsian groups. The Möbius maps defining \widehat{A}_{Γ} are also marked, where \widehat{g}_i is the image of g_i under the representation $\Gamma_0 \to \Gamma$.

conjugates the Bowen-Series (respectively, higher Bowen-Series) map of Γ_0 to a self-map of $\Lambda(\Gamma)$ that is orbit equivalent to Γ . These Kleinian groups arise out of pinching finitely many disjoint, simple, closed curves (on the surface \mathbb{D}/Γ_0) out of an explicit finite list. In particular, all such groups Γ are geometrically finite.

Let $\Gamma \in \partial \mathcal{B}(\Gamma_0)$ be a group that admits a Bowen-Series (respectively, higher Bowen-Series) map A_{Γ} . Then the canonical extension $\widehat{A}_{\Gamma} : K(\Gamma) \setminus int R_{\Gamma} \to K(\Gamma)$ can be conformally mated with polynomials lying in the principal hyperbolic component of degree 2k - 3 (respectively, $(k - 1)^2$).

We refer the reader to [MM21, Remark 7.20] for a precise definition of conformal mateability of canonical extensions of the Bowen-Series/ higher Bowen-Series maps with polynomials lying in principal hyperbolic components (the definition is analogous to Definition 30).

The finiteness part of Theorem 50 underscores the incompatibility between group invariant geodesic laminations and polynomial laminations (see [Kiw04] for details on polynomial laminations) by establishing that the equivalence relation on \mathbb{S}^1 induced by a group invariant geodesic lamination on \mathbb{D} is seldom invariant under A_{Γ_0} (since $A_{\Gamma_0}|_{\mathbb{S}^1}$ is topologically conjugate to $z^d|_{\mathbb{S}^1}$ for some $d \ge 2$, invariance under A_{Γ_0} should be thought of as z^d -invariance).

The proof of existence of a conformal mating between $\widehat{A}_{\Gamma} : K(\Gamma) \setminus \operatorname{int} R_{\Gamma} \to K(\Gamma)$ and polynomials lying in principal hyperbolic components has two main steps. The first one is to topologically realize the action of $A_{\Gamma}|_{\Lambda_{\Gamma}}$ by the dynamics of a

postcritically finite polynomial P_{Γ} on its Julia set, which is the content of [MM21, Theorem 7.16]. Once this is achieved, one needs to replace the dynamics of P_{Γ} on periodic Fatou components by the action of Bowen-Series/ higher Bowen-Series maps of suitable punctured sphere Fuchsian groups. This involves a rather delicate surgery technique using David homeomorphisms.

References

- AFW15. Matthias Aschenbrenner, Stefan Friedl, and Henry Wilton. *3-manifold groups*. EMS Series of Lectures in Mathematics. European Mathematical Society (EMS), Zürich, 2015.
- AGM16. Ian Agol, Daniel Groves, and Jason Fox Manning. An alternate proof of Wise's malnormal special quotient theorem. *Forum Math. Pi*, 4:e1, 54, 2016.
- Ago13. Ian Agol. The virtual Haken conjecture. *Doc. Math.*, 18:1045–1087, 2013. With an appendix by Agol, Daniel Groves, and Jason Manning.
- Alio5. Emina Alibegović. A combination theorem for relatively hyperbolic groups. Bull. London Math. Soc., 37(3):459–466, 2005.
- AR16. Magnus Aspenberg and Pascale Roesch. Newton maps as matings of cubic polynomials. Proc. Lond. Math. Soc. (3), 113(1):77–112, 2016.
- AS76. D. Aharonov and H. S. Shapiro. Domains on which analytic functions satisfy quadrature identities. J. Analyse Math., 30:39–73, 1976.
- BD18. Laurent Bartholdi and Dzmitry Dudko. Algorithmic aspects of branched coverings IV/V. Expanding maps. *Trans. Amer. Math. Soc.*, 370(11):7679–7714, 2018.
- BEK12a. Xavier Buff, Adam L. Epstein, and Sarah Koch. Twisted matings and equipotential gluings. Ann. Fac. Sci. Toulouse Math. (6), 21(5):995–1031, 2012.
- BEK⁺12b. Xavier Buff, Adam L. Epstein, Sarah Koch, Daniel Meyer, Kevin Pilgrim, Mary Rees, and Tan Lei. Questions about polynomial matings. *Ann. Fac. Sci. Toulouse Math.* (6), 21(5):1149–1176, 2012.
- Ber60. Lipman Bers. Simultaneous uniformization. Bull. Amer. Math. Soc., 66:94-97, 1960.
- Ber61. Lipman Bers. Uniformization by Beltrami equations. Comm. Pure Appl. Math., 14:215– 228, 1961.
- Bes02. M. Bestvina. R-trees in topology, geometry, and group theory. Handbook of geometric topology, Amsterdam: North-Holland, pages 55–91, 2002.
- BF92. M. Bestvina and M. Feighn. A Combination theorem for Negatively Curved Groups. J. Diff. Geom., vol 35, pages 85–101, 1992.
- BM17. Mario Bonk and Daniel Meyer. *Expanding Thurston maps*, volume 225 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2017.
- Bow79. R Bowen. Hausdorff dimension of quasicircles. Inst. Hautes Études Sci. Publ. Math., 50:11–25, 1979.
- Bow07. B. H. Bowditch. The Cannon-Thurston map for punctured surface groups. *Math. Z. 255*, pages 35–76, (2007).
- Bow12. B. H. Bowditch. Relatively hyperbolic groups. Internat. J. Algebra and Computation. 22, 1250016, 66pp, 2012.
- BP94. S. Bullett and C. Penrose. Mating quadratic maps with the modular group. *Invent. Math.*, 115:483–511, 1994.
- Bro65. Hans Brolin. Invariant sets under iteration of rational functions. *Ark. Mat.*, 6:103–144 (1965), 1965.
- BS79. R. Bowen and C. Series. Markov maps associated with Fuchsian groups. Inst. Hautes Études Sci. Publ. Math., 50:153–170, 1979.
- BW12. Nicolas Bergeron and Daniel T. Wise. A boundary criterion for cubulation. Amer. J. Math., 134(3):843–859, 2012.
- CG93. L. Carleson and T. W. Gamelin. Complex Dynamics. Springer, Berlin, 1993.

- Ché. Arnaud Chéritat. Polynomial matings on the riemann sphere. https://www.math. univ-toulouse.fr/~cheritat/MatMovies/.
- Ché12. Arnaud Chéritat. Tan Lei and Shishikura's example of non-mateable degree 3 polynomials without a Levy cycle. *Ann. Fac. Sci. Toulouse Math.* (6), 21(5):935–980, 2012.
- CPT12. Guizhen Cui, Wenjuan Peng, and Lei Tan. On a theorem of Rees-Shishikura. *Ann. Fac. Sci. Toulouse Math.* (6), 21(5):981–993, 2012.
- CR80. E. M. Coven and W. L. Reddy. Positively expansive maps of compact manifolds. In *Global theory of dynamical systems (Proc. Internat. Conf., Northwestern Univ., Evanston, Ill., 1979)*, volume 819 of *Lecture Notes in Math.*, pages 96–110. Springer, Berlin, 1980.
- CT07. J. Cannon and W. P. Thurston. Group Invariant Peano Curves. *Geom. Topol. 11*, pages 1315–1355, 2007.
- Dah03. François Dahmani. Combination of convergence groups. Geom. Topol., 7:933-963, 2003.
- DeZ20. Alexandre DeZotti. Some connections of complex dynamics. https://arxiv.org/ abs/2006.16386, 2020.
- DH84. A. Douady and J. H. Hubbard. Étude dynamique des polynômes complexes. Partie I, volume 84 of Publications Mathématiques d'Orsay [Mathematical Publications of Orsay]. Université de Paris-Sud, Département de Mathématiques, Orsay, 1984.
- DH85. A. Douady and J. H. Hubbard. Étude dynamique des polynômes complexes. Partie II, volume 85 of Publications Mathématiques d'Orsay [Mathematical Publications of Orsay]. Université de Paris-Sud, Département de Mathématiques, Orsay, 1985. With the collaboration of P. Lavaurs, Tan Lei and P. Sentenac.
- DH93. A. Douady and J. H. Hubbard. A proof of thurston's topological characterization of rational functions. *Acta Math.*, 171:263–297, 1993.
- DKL14. S. Dowdall, R. P. Kent, and C. J. Leininger. Pseudo-anosov subgroups of fibered 3manifold groups. Groups Geom. Dyn. 8, no. 4, pages 1247–1282, 2014.
- DM16. S. Das and M. Mj. Semiconjugacies Between Relatively Hyperbolic Boundaries. Groups Geom. Dyn. 10, pages 733–752, 2016.
- Dou83. A. Douady. Systèmes dynamiques holomorphes. In Séminaire Bourbaki, volume 1982/83, pages 39–63. Astérisque, 105–106, Soc. Math. France, Paris, 1983.
- Dur83. Peter L. Duren. Univalent functions, volume 259 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, New York, 1983.
- ES12. Adam Epstein and Thomas Sharland. A classification of bicritical rational maps with a pair of period two superattracting cycles. *Ann. Fac. Sci. Toulouse Math.* (6), 21(5):907–934, 2012.
- Far98. B. Farb. Relatively hyperbolic groups. Geom. Funct. Anal. 8, pages 810–840, 1998.
- Fat19. P. Fatou. Sur les équations fonctionnelles. Bull. Soc. Math. France, 47:161-271, 1919.
- Fat20a. P. Fatou. Sur les équations fonctionnelles. Bull. Soc. Math. France, 48:33-94, 1920.
- Fat20b. P. Fatou. Sur les équations fonctionnelles. Bull. Soc. Math. France, 48:208-314, 1920.
- Fat26. P. Fatou. Sur l'itération des fonctions transcendantes Entières. Acta Math., 47(4):337–370, 1926.
- Fat29. P. Fatou. Notice sur les travaux scientifiques de M. P. Fatou. Paris, 1929.
- FLP79. A. Fathi, M. Laudenbach, and V. Poenaru. Travaux de Thurston sur les surfaces. Asterisque 66-67, pages 1–284, 1979.
- Gau16. F. Gautero. Geodesics in Trees of Hyperbolic and Relatively Hyperbolic Spaces. Proc. Edinburgh. Math. Soc. 59(3), pages 701–740, 2016.
- Ger98. S. M. Gersten. Cohomological lower bounds for isoperimetric functions on groups. *Topology*, 37(5):1031–1072, 1998.
- GH09. François Gautero and Michael Heusener. Cohomological characterization of relative hyperbolicity and combination theorem. *Publ. Mat.*, 53(2):489–514, 2009.
- Gro85. M. Gromov. Hyperbolic Groups. in Essays in Group Theory, ed. Gersten, MSRI Publ., vol.8, Springer Verlag, pages 75–263, 1985.
- GS20. Subhojoy Gupta and Harish Seshadri. Complex geometry of Teichmüller domains. In Handbook of Teichmüller theory. Vol. VII, volume 30 of IRMA Lect. Math. Theor. Phys., pages 63–88. Eur. Math. Soc., Zürich, 2020.

- GW11. F. Gautero and R. Weidmann. An algebraic combination theorem for graphs of relatively hyperbolic groups. *preprint*, *University of Nice*, 2011.
- Ham05. U. Hamenstaedt. Word hyperbolic extensions of surface groups . preprint, arXiv:math/0505244, 2005.
- Hem04. John Hempel. 3-manifolds. AMS Chelsea Publishing, Providence, RI, 2004. Reprint of the 1976 original.
- Hru10. G. Christopher Hruska. Relative hyperbolicity and relative quasiconvexity for countable groups. Algebr. Geom. Topol., 10(3):1807–1856, 2010.
- HT04. Peter Haïssinsky and Lei Tan. Convergence of pinching deformations and matings of geometrically finite polynomials. *Fund. Math.*, 181(2):143–188, 2004.
- Hub06. John Hamal Hubbard. Teichmüller theory and applications to geometry, topology, and dynamics. Vol. 1. Matrix Editions, Ithaca, NY, 2006. Teichmüller theory, With contributions by Adrien Douady, William Dunbar, Roland Roeder, Sylvain Bonnot, David Brown, Allen Hatcher, Chris Hruska and Sudeb Mitra, With forewords by William Thurston and Clifford Earle.
- HW08. F. Haglund and D. T. Wise. Special cube complexes. *Geom. Funct. Anal. 17 no. 5*, pages 1551–1620, 2008.
- HW10. Tim Hsu and Daniel T. Wise. Cubulating graphs of free groups with cyclic edge groups. Amer. J. Math., 132(5):1153–1188, 2010.
- HW12. F. Haglund and D. T. Wise. A combination theorem for special cube complexes. Ann. of Math. (2) 176, no. 3, pages 1427–1482, 2012.
- HW15a. Mark F. Hagen and Daniel T. Wise. Cubulating hyperbolic free-by-cyclic groups: the general case. *Geom. Funct. Anal.*, 25(1):134–179, 2015.
- HW15b. Tim Hsu and Daniel T. Wise. Cubulating malnormal amalgams. *Invent. Math.*, 199(2):293–331, 2015.
- HW16. Mark F. Hagen and Daniel T. Wise. Cubulating hyperbolic free-by-cyclic groups: the irreducible case. Duke Math. J., 165(9):1753–1813, 2016.
- Jul18. Gaston Julia. Mémoire sur l'itération des fonctions rationnelles. Journal de Mathématiques Pures et Appliquées, 1:47–246, 1918.
- Jul22. Gaston Julia. Mémoire sur la permutabilité des fractions rationnelles. Ann. Sci. École Norm. Sup. (3), 39:131–215, 1922.
- Kabl6. Y. Kabaya. Deformation of quasi-fuchsian groups. https://www.math.kyoto-u. ac.jp/~kabaya/galleries/quasifuchsian_deformation/quasifuchsian_ deformation.html, 2016.
- Kap01a. Ilya Kapovich. The combination theorem and quasiconvexity. Internat. J. Algebra Comput., 11(2):185–216, 2001.
- Kap01b. Michael Kapovich. Hyperbolic manifolds and discrete groups, volume 183 of Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 2001.
- Kiw04. J. Kiwi. Real laminations and the topological dynamics of complex polynomials. Adv. Math., 184(2):207–267, 2004.
- Kle83. Felix Klein. Neue Beiträge zur Riemann'schen Functionentheorie. Math. Ann., 21(2):141– 218, 1883.
- KM12. Jeremy Kahn and Vladimir Markovic. Immersing almost geodesic surfaces in a closed hyperbolic three manifold. *Ann. of Math.* (2), 175(3):1127–1190, 2012.
- KS21. M. Kapovich and P. Sardar. Trees of hyperbolic spaces. book, in preparation, 2021.
- LLM20. Russell Lodge, Yusheng Luo, and Sabyasachi Mukherjee. Circle packings, kissing reflection groups and critically fixed anti-rational maps. arXiv:2007.03558, 2020.
- LLMM18a. S.-Y. Lee, M. Lyubich, N. G. Makarov, and S. Mukherjee. Dynamics of Schwarz reflections: the mating phenomena. https://arxiv.org/abs/1811.04979, 2018.
- LLMM18b. S.-Y. Lee, M. Lyubich, N. G. Makarov, and S. Mukherjee. Schwarz reflections and the Tricorn. https://arxiv.org/abs/1812.01573, 2018.
- LLMM19. S.-Y. Lee, M. Lyubich, N. G. Makarov, and S. Mukherjee. Schwarz reflections and anti-holomorphic correspondences. https://arxiv.org/abs/1907.09107, 2019.
- LM97. M. Lyubich and Y. Minsky. Laminations in holomorphic dynamics. J. Differential Geom., 47:17–94, 1997.

- LM14. S.-Y. Lee and N. Makarov. Sharpness of connectivity bounds for quadrature domains. https://arxiv.org/abs/1411.3415, November 2014.
- LM16. Seung-Yeop Lee and Nikolai G. Makarov. Topology of quadrature domains. J. Amer. Math. Soc., 29(2):333–369, 2016.
- LMM19. K. Lazebnik, N. G. Makarov, and S. Mukherjee. Univalent polynomials and Hubbard trees. https://arxiv.org/abs/1908.05813, 2019.
- LMM20. K. Lazebnik, N. G. Makarov, and S. Mukherjee. Bers slices in families of univalent maps. https://arxiv.org/abs/2007.02429, 2020.
- LMMN20. M. Lyubich, S. Merenkov, S. Mukherjee, and D. Ntalampekos. David extension of circle homeomorphisms, welding, mating, and removability. https://arxiv.org/abs/ 2010.11256, 2020.
- LOW09. Liulan Li, Ken'ichi Ohshika, and Xiantao Wang. On Klein-Maskit combination theorem in space. I. Osaka J. Math., 46(4):1097–1141, 2009.
- LOW15. Liulan Li, Ken'ichi Ohshika, and Xiantao Wang. On Klein-Maskit combination theorem in space II. Kodai Math. J., 38(1):1–22, 2015.
- Luo19. Yusheng Luo. Limits of rational maps, R-trees and barycentric extension. https:// arxiv.org/abs/1905.00915, 2019.
- Luo21a. Yusheng Luo. On geometrically finite degenerations I: boundaries of main hyperbolic components. https://arxiv.org/abs/2101.07880, 2021.
- Luo21b. Yusheng Luo. On geometrically finite degenerations II: convergence and divergence. https://arxiv.org/abs/2102.00357, 2021.
- Lyu17. M. Lyubich. *Conformal Geometry and Dynamics of Quadratic Polynomials, vol I-II.* in preparation, 2017. http://www.math.stonybrook.edu/~mlyubich/book.pdf.
- Mas65a. Bernard Maskit. Construction of Kleinian groups. In Proc. Conf. Complex Analysis (Minneapolis, 1964), pages 281–296. Springer, Berlin, 1965.
- Mas65b. Bernard Maskit. On Klein's combination theorem. *Trans. Amer. Math. Soc.*, 120:499–509, 1965.
- Mas68. Bernard Maskit. On Klein's combination theorem. II. *Trans. Amer. Math. Soc.*, 131:32–39, 1968.
- Mas71. Bernard Maskit. On Klein's combination theorem. III. In Advances in the Theory of Riemann Surfaces (Proc. Conf., Stony Brook, N.Y., 1969), Ann. of Math. Studies, No. 66, pages 297–316. Princeton Univ. Press, Princeton, N.J., 1971.
- Mas80. B. Maskit. Kleinian Groups. Springer Grundlehren 287, 1980.
- Mas93. Bernard Maskit. On Klein's combination theorem. IV. Trans. Amer. Math. Soc., 336(1):265–294, 1993.
- McM90. C. McMullen. Iteration on Teichmuller Space. Invent. math., 9:425-454, 1990.
- McM98. C. T. McMullen. Renormalization and 3-Manifolds which Fiber over the Circle. *Princeton University Press*, 1998.
- Mey09. Daniel Meyer. Expanding thurston maps as quotients. https://arxiv.org/abs/0910. 2003, 2009.
- Mey14. Daniel Meyer. Unmating of rational maps: sufficient criteria and examples. In Frontiers in complex dynamics, volume 51 of Princeton Math. Ser., pages 197–233. Princeton Univ. Press, Princeton, NJ, 2014.
- Mil00. John Milnor. Periodic orbits, externals rays and the Mandelbrot set: an expository account. In Géométrie complexe et systèmes dynamiques (Orsay, 1995), volume 261, pages xiii, 277– 333. Astérisque, 2000.
- Mil04. John Milnor. Pasting together Julia sets: a worked out example of mating. *Experiment*. *Math.*, 13(1):55–92, 2004.
- Mil06. J. Milnor. Dynamics in one complex variable, volume 160 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, third edition, 2006.
- Mit98a. M. Mitra. Cannon-Thurston Maps for Hyperbolic Group Extensions. *Topology 37*, pages 527–538, 1998.
- Mit98b. M. Mitra. Cannon-Thurston Maps for Trees of Hyperbolic Metric Spaces. Jour. Diff. Geom.48, pages 135–164, 1998.

- Mj14a. M. Mj. Cannon-Thurston Maps for Surface Groups. Ann. of Math., 179(1), pages 1–80, 2014.
- Mj14b. M. Mj. Ending Laminations and Cannon-Thurston Maps, with an appendix by S. Das and M. Mj. Geom. Funct. Anal. 24, pages 297–321, 2014.
- Mj17. M. Mj. Cannon-Thurston Maps for Kleinian Groups. Forum Math. Pi 5, e1, 49 pp., 2017.
- Mj20. M. Mj. Tight trees and model geometries of surface bundles over graphs. J. Lond. Math. Soc. (2) 102, no. 3, pages 1178–1222, 2020.
- MM21. M. Mj and S. Mukherjee. Combining rational maps and kleinian groups via orbit equivalence. https://arxiv.org/abs/2103.00835, 2021.
- MMS19. Jason Manning, Mahan Mj, and Michah Sageev. Cubulating Surface-by-free Groups. preprint, arxiv:1908.03545, page 46, 2019.
- MO21. A. Martin and D. Osajda. A combination theorem for combinatorially non-positively curved complexes of hyperbolic groups. *Mathematical Proceedings of the Cambridge Philosophical Society*, 2021.
- MR08. M. Mj and L. Reeves. A Combination Theorem for Strong Relative Hyperbolicity. *Geom. Topol.* 12, pages 1777 1798, 2008.
- MS84. John W. Morgan and Peter B. Shalen. Valuations, trees, and degenerations of hyperbolic structures. I. Ann. of Math. (2), 120(3):401–476, 1984.
- MS88a. John W. Morgan and Peter B. Shalen. Degenerations of hyperbolic structures. II. Measured laminations in 3-manifolds. *Ann. of Math.* (2), 127(2):403–456, 1988.
- MS88b. John W. Morgan and Peter B. Shalen. Degenerations of hyperbolic structures. III. Actions of 3-manifold groups on trees and Thurston's compactness theorem. *Ann. of Math.* (2), 127(3):457–519, 1988.
- MS12. M. Mj and P. Sardar. A combination theorem for metric bundles. *Geom. Funct. Anal. 22, no.* 6, pages 1636–1707, 2012.
- MT12. Inna Mashanova and Vladlen Timorin. Captures, matings and regluings. *Ann. Fac. Sci. Toulouse Math.* (6), 21(5):877–906, 2012.
- Ota98. J. P. Otal. Thurston's hyperbolization of Haken manifolds. *Surveys in differential geometry, Vol. III (Cambridge, MA, 1996)*, pages 77–194, 1998.
- Ota01. Jean-Pierre Otal. The hyperbolization theorem for fibered 3-manifolds, volume 7 of SMF/AMS Texts and Monographs. American Mathematical Society, Providence, RI; Société Mathématique de France, Paris, 2001. Translated from the 1996 French original by Leslie D. Kay.
- Pil03. K. M. Pilgrim. Combinations of complex dynamical systems. Lecture Notes in Mathematics. Springer, 2003.
- PL98. Kevin M. Pilgrim and Tan Lei. Combining rational maps and controlling obstructions. Ergodic Theory Dynam. Systems, 18(1):221–245, 1998.
- PM12. Carsten Lunde Petersen and Daniel Meyer. On the notions of mating. Ann. Fac. Sci. Toulouse Math. (6), 21(5):839–876, 2012.
- PW18. Piotr Przytycki and Daniel T. Wise. Mixed 3-manifolds are virtually special. J. Amer. Math. Soc., 31(2):319–347, 2018.
- Ree92. Mary Rees. A partial description of parameter space of rational maps of degree two. I. Acta Math., 168(1-2):11–87, 1992.
- Ree10. Mary Rees. Multiple equivalent matings with the aeroplane polynomial. Ergodic Theory Dynam. Systems, 30(4):1239–1257, 2010.
- Ree16. Mary Rees. One hundred years of complex dynamics. Proc. A., 472(2185):20150453, 16, 2016.
- Roe12. Pascale Roesch. Introduction [Workshop on Polynomial Matings]. Ann. Fac. Sci. Toulouse Math. (6), 21(5):iii–x, 2012. Held in Toulouse, June 8–11, 2011.
- Sag95. Michah Sageev. Ends of group pairs and non-positively curved cube complexes. Proc. London Math. Soc. (3), 71(3):585–617, 1995.
- Sak91. Makoto Sakai. Regularity of a boundary having a Schwarz function. Acta Math., 166(3-4):263–297, 1991.
- Sel97. Z. Sela. Acylindrical accessibility for groups. Invent. Math., 129(3):527-565, 1997.

- Sha19. Thomas Sharland. Matings of cubic polynomials with a fixed critical point, Part I: Thurston obstructions. Conform. Geom. Dyn., 23:205–220, 2019.
- Shi00. Mitsuhiro Shishikura. On a theorem of M. Rees for matings of polynomials. In *The Mandelbrot set, theme and variations*, volume 274 of *London Math. Soc. Lecture Note Ser.*, pages 289–305. Cambridge Univ. Press, Cambridge, 2000.
- Sie42. Carl Ludwig Siegel. Iteration of analytic functions. Ann. of Math. (2), 43:607-612, 1942.
- Sko96. Richard K. Skora. Splittings of surfaces. J. Amer. Math. Soc., 9(2):605-616, 1996.
- SL00. Mitsuhiro Shishikura and Tan Lei. A family of cubic rational maps and matings of cubic polynomials. *Experiment. Math.*, 9(1):29–53, 2000.
- SM98. D. Sullivan and C. T. McMullen. Quasiconformal homeomorphisms and dynamics III: The Teichmüller space of a holomorphic dynamical system. *Adv. Math.*, 135:351–395, 1998.
- Sul85. D. Sullivan. Quasiconformal Homeomorphisms and Dynamics I: Solution of the Fatou-Julia problem on Wandering Domains. *Annals of Math. 122*, pages 401–418, 1985.
- SW79. P. Scott and C. T. C. Wall. Topological Methods in Group Theory. Homological group theory (C. T. C. Wall, ed.), London Math. Soc. Lecture Notes Series, vol. 36, Cambridge Univ. Press, 1979.
- Tan92. Lei Tan. Matings of quadratic polynomials. Ergodic Theory Dynam. Systems, 12(3):589– 620, 1992.
- Tan97. Lei Tan. Branched coverings and cubic Newton maps. Fund. Math., 154(3):207–260, 1997.
- Thu80. W. P. Thurston. The Geometry and Topology of 3-Manifolds. *Princeton University Notes*, 1980.
- Thu86a. W. P. Thurston. Hyperbolic Structures on 3-Manifolds,I: Deformations of acylindrical manifolds. Annals of Math.124, pages 203–246, 1986.
- Thu86b. W. P. Thurston. Hyperbolic Structures on 3-Manifolds, II: Surface Groups and 3-Manifolds which Fiber over the Circle. *preprint*, arXiv:math.GT/9801045, 1986.
- Thu86c. W. P. Thurston. Hyperbolic Structures on 3-Manifolds,III: Deformations of 3-manifolds with incompressible boundary. *preprint*, 1986.
- VS93. È. B. Vinberg and O. V. Shvartsman. Discrete groups of motions of spaces of constant curvature. In *Geometry*, II, volume 29 of *Encyclopaedia Math. Sci.*, pages 139–248. Springer, Berlin, 1993.
- Wis02. Daniel T. Wise. The residual finiteness of negatively curved polygons of finite groups. Invent. Math., 149(3):579–617, 2002.
- Wis12. Daniel T. Wise. From riches to raags: 3-manifolds, right-angled Artin groups, and cubical geometry, volume 117 of CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2012.
- Wis21. D. T. Wise. The structure of groups with a quasi-convex hierarchy. Annals of Math. Studies AMS 209, 2021.
- YZ01. Michael Yampolsky and Saeed Zakeri. Mating Siegel quadratic polynomials. J. Amer. Math. Soc., 14(1):25–78, 2001.